

## RESEARCH ARTICLE

WILEY

# Inverse nodal problem for Dirac operator with integral type nonlocal boundary conditions

A. Sinan Ozkan<sup>1</sup>  | İbrahim Adalar<sup>2</sup> 

<sup>1</sup>Department of Mathematics, Faculty of Science, Sivas Cumhuriyet University, Sivas, Turkey

<sup>2</sup>Zara Veysel Dursun Colleges of Applied Sciences, Sivas Cumhuriyet University, Sivas, Turkey

## Correspondence

A. Sinan Ozkan, Department of Mathematics, Faculty of Science, Sivas Cumhuriyet University 58140 Sivas, Sivas, Turkey.

Email: sozkan@cumhuriyet.edu.tr

Communicated by: S. Georgiev

In this paper, Dirac operator with some integral type nonlocal boundary conditions is studied. We show that the coefficients of the problem can be uniquely determined by a dense set of nodal points. Moreover, we give an algorithm for the reconstruction of some coefficients of the operator.

## KEYWORDS

Dirac operator, inverse nodal problem, nonlocal boundary condition

## MSC CLASSIFICATION

34A55, 34L05, 34K29, 34K10

## 1 | INTRODUCTION

The inverse nodal problem was posed and solved firstly by McLaughlin for a Sturm–Liouville operator.<sup>1</sup> McLaughlin showed that the potential of a Sturm–Liouville problem can be determined by a given dense subset of nodal points. Later on, Hald and McLaughlin give some numerical schemes for the reconstruction of the potential.<sup>2</sup> X.F. Yang gave an algorithm for the solution of the inverse nodal Sturm–Liouville problem.<sup>3</sup> His method is the first of the algorithms still in use today. Inverse nodal problems for various Sturm–Liouville operators have been studied in the papers (<sup>4–23</sup>).

Nonlocal boundary conditions appear when we cannot measure data directly at the boundary. These kinds of conditions arise in various some applied problems of biology, biotechnology, physics, and so on. Two types of nonlocal boundary conditions come to the fore. One class of them is called integral type conditions, and the other is the Bitsadze–Samarskii-type conditions. Some inverse problems for a class of Sturm–Liouville operators with nonlocal boundary conditions are investigated in earlier studies.<sup>24,25</sup> In particular, inverse nodal problems for this-type operators with different nonlocal integral boundary conditions are studied in previous research<sup>26–29</sup>. The inverse nodal problems for Dirac operators with local and separated boundary conditions are studied in earlier studies<sup>30–36</sup>. In their works, it is shown that the zeros of the first components of the eigenfunctions determine the coefficients of operator. The uniqueness of the recovery of the Dirac operator with nonseparated boundary conditions is investigated in earlier works<sup>37–39</sup>. In Yang and Yurko,<sup>40</sup> nonlocal conditions together with the Dirac system are considered. They give some uniqueness theorems according to the classical spectral data. As far as we know, inverse nodal problem for the Dirac system with nonlocal boundary conditions has not been considered before. In the present paper, we consider Dirac operator under some integral type nonlocal boundary conditions and obtain the uniqueness of coefficients of the problem according to a set of nodal points. Moreover, we give an algorithm for the reconstruction of these coefficients. We consider the boundary value problem  $L$  generated by the system of Dirac differential equations system:

$$\ell [Y(x)] := BY'(x) + Q(x)Y(x) = \lambda Y(x), x \in (0, \pi), \quad (1)$$

subject to the boundary conditions

$$U(y) := y_1(0) \sin \alpha + y_2(0) \cos \alpha - I_f(Y) = 0, \quad (2)$$

$$V(y) := y_1(\pi) \sin \beta + y_2(\pi) \cos \beta - I_g(Y) = 0, \quad (3)$$

where  $B = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ ,  $Q(x) = \begin{pmatrix} V(x) + m & 0 \\ 0 & V(x) - m \end{pmatrix}$ ,  $Y(x) = \begin{pmatrix} y_1(x) \\ y_2(x) \end{pmatrix}$ ,  $I_f(Y) = \int_0^\pi (f_1(x)y_1(x) + f_2(x)y_2(x)) dx$ ,  $I_g(Y) = \int_0^\pi (g_1(x)y_1(x) + g_2(x)y_2(x)) dx$ ,  $\alpha$ ,  $\beta$ , and  $m$  are real constants, and  $\lambda$  is the spectral parameter. We assume that  $V(x)$ ,  $f_i(x)$  and  $g_i(x)$  are real-valued functions in the class of  $W_2^1(0, \pi)$  for  $i = 1, 2$  and put  $F_0 := f_1(0) \sin \alpha + f_2(0) \cos \alpha$ ,  $F_\pi := f_1(\pi) \sin \beta + f_2(\pi) \cos \beta$ ,  $G_0 := g_1(0) \sin \alpha + g_2(0) \cos \alpha$  and  $G_\pi := g_1(\pi) \sin \beta + g_2(\pi) \cos \beta$ .

## 2 | SPECTRAL PROPERTIES OF THE PROBLEM

Let  $S(x, \lambda) = \begin{pmatrix} S_1(x, \lambda) \\ S_2(x, \lambda) \end{pmatrix}$  and  $C(x, \lambda) = \begin{pmatrix} C_1(x, \lambda) \\ C_2(x, \lambda) \end{pmatrix}$  be the solutions of (1) under the initial conditions

$$S_1(0, \lambda) = 0, \quad S_2(0, \lambda) = -1,$$

$$C_1(0, \lambda) = 1, \quad C_2(0, \lambda) = 0,$$

respectively.  $C(x, \lambda)$  and  $S(x, \lambda)$  are entire functions of  $\lambda$  for each fixed  $x$  and the following asymptotic relations hold for  $|\lambda| \rightarrow \infty$  (see Keskin and Ozkan<sup>30</sup>).

$$C_1(x, \lambda) = \cos(\lambda x - \omega(x)) + \frac{m^2 x}{2\lambda} \sin(\lambda x - \omega(x)) + o\left(\frac{e^{|\tau|x}}{\lambda}\right), \quad (4)$$

$$C_2(x, \lambda) = \sin(\lambda x - \omega(x)) - \frac{m}{\lambda} \sin(\lambda x - \omega(x)) - \frac{m^2 x}{2\lambda} \cos(\lambda x - \omega(x)) + o\left(\frac{e^{|\tau|x}}{\lambda}\right), \quad (5)$$

$$S_1(x, \lambda) = \sin(\lambda x - \omega(x)) + \frac{m}{\lambda} \sin(\lambda x - \omega(x)) - \frac{m^2 x}{2\lambda} \cos(\lambda x - \omega(x)) + o\left(\frac{e^{|\tau|x}}{\lambda}\right), \quad (6)$$

$$S_2(x, \lambda) = -\cos(\lambda x - \omega(x)) - \frac{m^2 x}{2\lambda} \sin(\lambda x - \omega(x)) + o\left(\frac{e^{|\tau|x}}{\lambda}\right), \quad (7)$$

where  $\omega(x) = \int_0^x V(t)dt$  and  $\tau = \text{Im}\lambda$ . The characteristic function of problem (1)–(3) is

$$\Delta(\lambda) = \det \begin{pmatrix} U(C) & U(S) \\ V(C) & V(S) \end{pmatrix} \quad (8)$$

and the zeros of  $\Delta(\lambda)$  coincide with the eigenvalues of problems (1)–(3). Clearly,  $\Delta(\lambda)$  is entire function and so the problem has a discrete spectrum. Using the asymptotic formulas (4)–(7) in (8), one can easily obtain

$$\begin{aligned} \Delta(\lambda) &= \sin(\lambda\pi - \omega(\pi) + \beta - \alpha) - \frac{m^2\pi}{2\lambda} \cos(\lambda\pi - \omega(\pi) + \beta - \alpha) \\ &\quad - \frac{1}{\lambda} (f_1(\pi) \sin \beta + f_2(\pi) \cos \beta - f_2(0) \cos(\lambda\pi - \omega(\pi) + \beta)) \\ &\quad - \frac{1}{\lambda} (g_1(\pi) \sin(\lambda\pi - \omega(\pi) - \alpha) - g_2(\pi) \cos(\lambda\pi - \omega(\pi) - \alpha)) \\ &\quad + \frac{1}{\lambda} (f_1(0) \sin(\lambda\pi - \omega(\pi) + \beta) - g_1(0) \sin \alpha - g_2(0) \cos \alpha) \\ &\quad - \frac{m}{\lambda} \sin(\lambda\pi - \omega(\pi) + \beta - \alpha) \cos(\beta + \alpha) \cos(\beta - \alpha) \\ &\quad + \frac{m}{\lambda} \cos(\lambda\pi - \omega(\pi) + \beta - \alpha) \cos(\beta + \alpha) \sin(\beta - \alpha) + o\left(\frac{e^{|\tau|\pi}}{\lambda}\right) \end{aligned}$$

for sufficiently large  $|\lambda|$ . Let  $\{\lambda_n : n = 0, \pm 1, \pm 2, \dots\}$  be the set of eigenvalues. Since  $\Delta(\lambda) = \sin(\lambda\pi - \omega(\pi) + \beta - \alpha) + O\left(\frac{e^{|\text{r}|\lambda}}{\lambda}\right)$ , the numbers  $\lambda_n$  are real for  $|n| \rightarrow \infty$  and if Rouché's theorem is used, the eigenvalues satisfy the following asymptotic formula

$$\lambda_n = n + \frac{\omega(\pi) - \beta + \alpha}{\pi} + O\left(\frac{1}{n}\right), |n| \rightarrow \infty.$$

Moreover, we can write the following equation.

$$\begin{aligned} \left(1 + O\left(\frac{1}{\lambda_n}\right)\right) \tan(\lambda_n\pi - \omega(\pi) + \beta - \alpha) &= \frac{1}{\lambda_n} \left[ \frac{m^2\pi}{2} - F_0 - G_\pi + m \cos(\beta + \alpha) \sin(\alpha - \beta) \right] \\ &+ \frac{F_\pi + G_0}{\lambda_n \cos(\lambda\pi - \omega(\pi) + \beta - \alpha)} + o\left(\frac{1}{\lambda_n}\right). \end{aligned}$$

Since  $\left(1 + O\left(\frac{1}{\lambda_n}\right)\right)^{-1} = 1 + O\left(\frac{1}{\lambda_n}\right)$ , this implies that

$$\tan(\lambda_n\pi - \omega(\pi) + \beta - \alpha) = \frac{A_1}{\lambda_n} + \frac{A_2}{\lambda_n \cos(\lambda\pi - \omega(\pi) + \beta - \alpha)} + o\left(\frac{1}{\lambda_n}\right) \quad (9)$$

for sufficiently large  $|n|$ , where

$$A_1 = \frac{m^2\pi}{2} + m \cos(\beta + \alpha) \sin(\alpha - \beta) - (F_0 + G_\pi),$$

$$A_2 = F_\pi + G_0.$$

From (9), we can see that

$$\lambda_n\pi - \omega(\pi) + \beta - \alpha = n\pi + O\left(\frac{1}{n}\right), |n| \rightarrow \infty.$$

Hence,

$$\begin{aligned} \frac{1}{\cos(\lambda_n\pi - \omega(\pi) + \beta - \alpha)} &= \frac{1}{\cos\left(n\pi + O\left(\frac{1}{n}\right)\right)} \\ &= (-1)^n \left(1 + o\left(\frac{1}{n}\right)\right), |n| \rightarrow \infty. \end{aligned} \quad (10)$$

Using (9) and (10) together, we obtain the following lemma.

**Lemma 1.** *The following asymptotic relation is valid for  $|n| \rightarrow \infty$ .*

$$\lambda_n = n + \frac{1}{\pi} \int_0^\pi V(t) dt + \frac{\alpha - \beta}{\pi} + \frac{A_1}{n\pi} + \frac{(-1)^n A_2}{n\pi} + o\left(\frac{1}{n}\right). \quad (11)$$

### 3 | NODAL POINTS

Let  $\varphi(x, \lambda_n) = \begin{pmatrix} \varphi_1(x, \lambda_n) \\ \varphi_2(x, \lambda_n) \end{pmatrix}$  be the eigenfunction of (1)–(3) corresponding to the eigenvalue  $\lambda_n$ . It is clear that

$$\varphi_i(x, \lambda_n) = U(S(x, \lambda_n))C_i(x, \lambda_n) - U(C(x, \lambda_n))S_i(x, \lambda_n), i = 1, 2. \quad (12)$$

From (12) and Lemma 1, we can give the following asymptotic formula for sufficiently large  $|n|$

$$\begin{aligned} \varphi_1(x, \lambda_n) = & -\cos(\lambda_n x - \omega(x) - \alpha) - \frac{m^2 x}{2\lambda_n} \sin(\lambda_n x - \omega(x) - \alpha) \\ & - \frac{m \sin 2\alpha}{2\lambda_n} \sin(\lambda_n x - \omega(x) - \alpha) - \frac{m \sin^2 \alpha}{\lambda_n} \cos(\lambda_n x - \omega(x) - \alpha) \\ & + \frac{(-1)^n f_1(\pi)}{\lambda_n} (\cos \beta \cos(\lambda_n x - \omega(x) - \alpha) - \sin \beta \sin(\lambda_n x - \omega(x) - \alpha)) \\ & - \frac{(-1)^n f_2(\pi)}{\lambda_n} (\cos \beta \sin(\lambda_n x - \omega(x) - \alpha) + \sin \beta \cos(\lambda_n x - \omega(x) - \alpha)) \\ & + \frac{f_1(0)}{\lambda_n} (\sin \alpha \sin(\lambda_n x - \omega(x) - \alpha) - \cos \alpha \cos(\lambda_n x - \omega(x) - \alpha)) \\ & + \frac{f_2(0)}{\lambda_n} (\cos \alpha \sin(\lambda_n x - \omega(x) - \alpha) + \sin \alpha \cos(\lambda_n x - \omega(x) - \alpha)) \\ & + o\left(\frac{e^{|\tau|x}}{\lambda}\right). \end{aligned}$$

**Lemma 2.** For sufficiently large positive  $n$ ,  $\varphi_1(x, \lambda_n)$  has exactly  $n$  zeros, namely, nodal points,  $\{x_n^j : j = \overline{0, n-1}\}$  in the interval  $(0, \pi)$ :

The numbers  $\{x_n^j\}$  satisfy the following asymptotic formula:

$$\begin{aligned} x_n^j = & \frac{(j+1/2)\pi}{n} + \frac{\omega(x_n^j) + \alpha}{n} - \frac{(j+1/2)}{n} \left( \frac{\omega(\pi) + \alpha - \beta}{n} \right) \\ & - \frac{\omega(\pi) + \alpha - \beta}{n^2 \pi} (\omega(x_n^j) + \alpha) + \frac{1}{n^2} \left( \frac{m^2 x_n^j}{2} + \frac{m \sin 2\alpha}{2} + (-1)^n F_\pi - F_0 \right) \\ & - \frac{(j+1/2)}{n} \left( \frac{A_1 + (-1)^n A_2}{n^2} \right) + o\left(\frac{1}{n^2}\right) \end{aligned} \tag{13}$$

for  $n \rightarrow \infty$ .

*Proof.* Let  $\kappa_{n,j} := \lambda_n x_n^j - \omega(x_n^j) - \alpha$ . From  $\varphi_1(x_n^j, \lambda_n) = 0$ , we can write for  $n \rightarrow \infty$

$$\begin{aligned} 0 = & -\cos \kappa_{n,j} - \frac{m \sin 2\alpha}{2\lambda_n} \sin \kappa_{n,j} - \frac{m \sin^2 \alpha}{\lambda_n} \cos \kappa_{n,j} - \frac{m^2 x_n^j}{2\lambda_n} \sin \kappa_{n,j} + \\ & + \frac{(-1)^n f_1(\pi)}{\lambda_n} (\cos \kappa_{n,j} \cos \beta - \sin \kappa_{n,j} \sin \beta) + \\ & - \frac{(-1)^n f_2(\pi)}{\lambda_n} (\sin \kappa_{n,j} \cos \beta + \cos \kappa_{n,j} \sin \beta) + \\ & + \frac{f_1(0)}{\lambda_n} (\sin \kappa_{n,j} \sin \alpha - \cos \kappa_{n,j} \cos \alpha) + \\ & + \frac{f_2(0)}{\lambda_n} (\sin \kappa_{n,j} \cos \alpha + \cos \kappa_{n,j} \sin \alpha) + o\left(\frac{e^{|\tau|\pi}}{\lambda_n}\right). \end{aligned}$$

It can be obtained that

$$\begin{aligned} \tan\left(\kappa_{n,j} - \frac{\pi}{2}\right) &= \frac{\frac{m^2 x_n^j}{2\lambda_n} + \frac{m \sin 2\alpha}{2\lambda_n} + \frac{(-1)^n F_\pi - F_0}{\lambda_n} + o\left(\frac{1}{\lambda_n}\right)}{1 + O\left(\frac{1}{\lambda_n}\right)} \\ &= \frac{m^2 x_n^j}{2\lambda_n} + \frac{m \sin 2\alpha}{2\lambda_n} + \frac{(-1)^n F_\pi - F_0}{\lambda_n} + o\left(\frac{1}{\lambda_n}\right). \end{aligned}$$

Taking into account Taylor's expansion formulas of the arctangent, we get

$$\kappa_{n,j} = \left(j + \frac{1}{2}\right) \pi + \frac{1}{\lambda_n} \left(\frac{m^2 x_n^j}{2} + \frac{m \sin 2\alpha}{2} + (-1)^n F_\pi - F_0\right) + o\left(\frac{1}{\lambda_n}\right).$$

It follows from the last equality that

$$x_n^j = \frac{\left(j + \frac{1}{2}\right) \pi + \omega(x_n^j) + \alpha}{\lambda_n} + \frac{1}{\lambda_n^2} \left(\frac{m^2 x_n^j}{2} + \frac{m \sin 2\alpha}{2} + (-1)^n F_\pi - F_0\right) + o\left(\frac{1}{\lambda_n^2}\right).$$

Finally, using the asymptotic formula

$$\lambda_n^{-1} = \frac{1}{n} \left\{ 1 - \frac{\omega(\pi) + \alpha - \beta}{n\pi} - \frac{A_1}{n^2\pi} - \frac{(-1)^n A_2}{n^2\pi} + o\left(\frac{1}{n^3}\right) \right\},$$

we can obtain our desired formula: (13). □

#### 4 | INVERSE NODAL PROBLEM

Let  $X$  be the set of nodal points and  $\omega(\pi) = 0$ . For each fixed  $x \in (0, \pi)$ , we can choose a sequence  $(x_n^{j(n)}) \subset X$  so that  $x_n^{j(n)}$  converges to  $x$ . Then the following limits are exist and finite:

$$\lim_{n \rightarrow \infty} n \left( x_n^{j(n)} - \frac{\left(j(n) + \frac{1}{2}\right) \pi}{n} \right) = \psi_1(x), \tag{14}$$

where

$$\psi_1(x) = \omega(x) + \frac{(\beta - \alpha)x}{\pi} + \alpha$$

and

$$\lim_{n \rightarrow \infty} n^2 \pi \left( x_n^{j(n)} - \frac{\left(j(n) + \frac{1}{2}\right) \pi + \left(\omega(x_n^{j(n)}) + \alpha\right)}{n} + \frac{(j(n) + 1/2)}{n} \left(\frac{\alpha - \beta}{n}\right) \right) = \psi_2(x), \tag{15}$$

where

$$\psi_2(x) = \begin{cases} \psi_2^+(x), & n \text{ is even} \\ \psi_2^-(x), & n \text{ is odd} \end{cases}$$

and

$$\begin{aligned} \psi_2^+(x) &= (\beta - \alpha)(\omega(x) + \alpha) - x(A_1 + A_2) + \pi \left(\frac{m^2 x}{2} + \frac{m \sin 2\alpha}{2} + F_\pi - F_0\right), \\ \psi_2^-(x) &= (\beta - \alpha)(\omega(x) + \alpha) - x(A_1 - A_2) + \pi \left(\frac{m^2 x}{2} + \frac{m \sin 2\alpha}{2} - F_\pi - F_0\right). \end{aligned}$$

Therefore, proof of the following theorem is clear.

**Theorem 1.** The given dense subset of nodal points  $X$  uniquely determines the potential  $V(x)$  a.e. on  $(0, \pi)$  and the coefficients  $\alpha$ ,  $\beta$ ,  $F_\pi$ , and  $G_0$  in the boundary conditions;  $m$ ,  $V(x)$ ,  $\alpha$ ,  $\beta$ ,  $F_\pi$ , and  $G_0$  can be reconstructed by the following formulae:

**Step-1:** For each fixed  $x \in (0, \pi)$ , choose a sequence  $(x_n^{j(n)}) \subset X$  such that  $\lim_{n \rightarrow \infty} x_n^{j(n)} = x$ ;

**Step-2:** Find the function  $\psi_1(x)$  and  $\psi_2(x)$  from (14) and (15) and calculate

$$\begin{aligned}\alpha &= \psi_1(0) \\ \beta &= \psi_1(\pi) \\ V(x) &= \psi_1'(x) + \frac{\alpha - \beta}{\pi} \\ F_\pi &= \frac{\psi_2^+(0) - \psi_2^-(0)}{2\pi} \\ G_0 &= \frac{\psi_2^-(\pi) - \psi_2^+(\pi)}{2\pi}.\end{aligned}$$

Additionally, if  $m$  is known, then  $F_0$  and  $G_\pi$  can be found by the following formulas

$$\begin{aligned}F_0 &= -\frac{\psi_2^+(0) + \psi_2^-(0)}{2} + \alpha(\beta - \alpha) + m\pi \frac{\sin 2\alpha}{2}, \\ G_\pi &= \frac{\psi_2^-(\pi) + \psi_2^+(\pi)}{2\pi} - \frac{\alpha(\beta - \alpha)}{\pi} - m \frac{\sin 2\beta}{2}.\end{aligned}$$

**Example 1.** Let  $\{x_n^j\} \subset X$  be a subset of nodal points of problems (1)–(3), and the sequence  $\{x_n^j\}$  satisfies the following asymptotics:

$$\begin{aligned}x_n^j &= \frac{(j+1/2)\pi}{n} + \frac{\sin \frac{(j+1/2)\pi}{n} + \frac{\pi}{6}}{n} + \frac{(j+1/2)\pi}{6n^2} \\ &+ \frac{1}{6n^2} \left( \sin \frac{(j+1/2)\pi}{n} + \frac{\pi}{6} \right) + \frac{1}{n^2} \left( \frac{3}{2} \left( \frac{(j+1/2)\pi}{n} + \frac{1}{2} \right) + (-1)^n 2\pi \right) \\ &- \frac{(j+1/2)}{n} \left( \frac{(2-\sqrt{3})\pi + (-1)^n 4\pi}{2n^2} \right) + o\left(\frac{1}{n^2}\right).\end{aligned}$$

It can be calculated from (14) and (15) that

$$\begin{aligned}\psi_1(x) &= \sin x + \frac{1}{6}(x + \pi), \\ \psi_2^+(x) &= \frac{\pi}{6} \left( \sin x + \frac{\pi}{6} \right) - \frac{\pi}{2} (6 - \sqrt{3})x + \pi \left( \frac{3}{2} \left( x + \frac{1}{2} \right) + 2\pi \right), \\ \psi_2^-(x) &= \frac{\pi}{6} \left( \sin x + \frac{\pi}{6} \right) + \frac{\pi}{2} (2 + \sqrt{3})x + \pi \left( \frac{3}{2} \left( x + \frac{1}{2} \right) - 2\pi \right).\end{aligned}$$

Therefore, it is obtained by using the algorithm in Theorem 1,

$$\begin{aligned}\alpha &= \frac{\pi}{6}, \\ \beta &= \frac{\pi}{3}, \\ V(x) &= \cos x, \\ F_\pi &= 2\pi, \\ G_0 &= 0.\end{aligned}$$

If  $m = \sqrt{3}$ , then

$$F_0 = 0,$$

$$G_\pi = \frac{(\sqrt{3} + 1)\pi}{2}.$$

## 5 | CONCLUSION AND RECOMMENDATION

Problem (1)–(3) have countably many eigenvalues that satisfy the asymptotic formula (11). For the sequence of nodal points of the problem, the asymptotic formula (13) holds. If  $\omega(\pi) = 0$ , a dense set of nodal points uniquely determines the coefficients  $V(x)$ ,  $\alpha$ ,  $\beta$  and the terms  $F_\pi$  and  $G_0$ . Also, if a nodal points-sequence that satisfies the asymptotic expression (13) is given, all of these coefficients and terms can be reconstructed by using the steps in Theorem 1. Our main result includes the full algorithm for the solution of the inverse nodal problem, therefore it would be useful for some problems in the applied sciences. After this, inverse nodal problems for a Dirac system with some different conditions (e.g., parameter-dependent nonlocal boundary conditions or transmission conditions) can be investigated.

## ACKNOWLEDGEMENTS

The authors would like to thank the referees for their valuable comments which helped to improve the manuscript. The authors also wish to state that there are no funders to report for this submission.

## CONFLICT OF INTEREST

This work does not have any conflict of interest.

## ORCID

A. Sinan Ozkan  <https://orcid.org/0000-0002-9703-8982>

İbrahim Adalar  <https://orcid.org/0000-0002-4224-0972>

## REFERENCES

- McLaughlin JR. Inverse spectral theory using nodal points as data—A uniqueness result. *J Differ Equ.* 1988;73(2):354-362.
- Hald OH, McLaughlin JR. Solution of inverse nodal problems. *Inverse Probl.* 1989;5(3):307-347.
- Yang XF. A solution of the inverse nodal problem. *Inverse Probl.* 1997;13(1):203-213.
- Yang XF. A new inverse nodal problem. *J Differ Equ.* 2001;169(2):633-653.
- Buterin SA, Shieh CT. Inverse nodal problem for differential pencils. *Appl Math Lett.* 2009;22(8):1240-1247.
- Buterin SA, Shieh CT. Incomplete inverse spectral and nodal problems for differential pencils. *RM.* 2012;62(1):167-179.
- Cheng YH, Law CK, Tsay J. Remarks on a new inverse nodal problem. *J Math Anal Appl.* 2000;248(1):145-155.
- Law CK, Tsay J. On the well-posedness of the inverse nodal problem. *Inverse Probl.* 2001;17(5):1493-1512.
- Law CK, Shen CL, Yang CF. The inverse nodal problem on the smoothness of the potential function. *Inverse Probl.* 1999;15(1):253.
- Law CK, Yang CF. Reconstructing the potential function and its derivatives using nodal data. *Inverse Probl.* 1998;14(2):299-312.
- Yang CF. Inverse nodal problems of discontinuous Sturm–Liouville operator. *J Differ Equ.* 2013;254(4):1992-2014.
- Yang CF, Yang XP. Inverse nodal problems for the Sturm–Liouville equation with polynomially dependent on the eigenparameter. *Inverse Probl Sci Eng.* 2011;19(7):951-961.
- Yang CF. Inverse nodal problems for the Sturm–Liouville operator with eigenparameter dependent boundary conditions. *Oper Matrices.* 2012;6(1):63-77.
- Ozkan AS, Keskin B. Inverse nodal problems for Sturm–Liouville equation with eigenparameter-dependent boundary and jump conditions. *Inverse Probl Sci Eng.* 2015;23(8):1306-1312.
- Shieh CT, Yurko VA. Inverse nodal and inverse spectral problems for discontinuous boundary value problems. *J Math Anal Appl.* 2008;347(1):266-272.
- Wang YP, Yurko VA. On the inverse nodal problems for discontinuous Sturm–Liouville operators. *J Differ Equ.* 2016;260(5):4086-4109.
- Yurko VA. Inverse nodal problems for Sturm–Liouville operators on star-type graphs. *J Inverse Ill-Posed Probl.* 2008;16:715-722.
- Yurko VA. *Inverse Spectral Problems for Differential Operators and Their Applications.* Amsterdam: Gordon and Breach; 2000.
- Shen CL, Shieh CT. An inverse nodal problem for vectorial Sturm–Liouville equations. *Inverse Probl.* 2000;16(2):349-356.
- Hu YT, Bondarenko NP, Yang CF. Traces and inverse nodal problem for Sturm–Liouville operators with frozen argument. *Appl Math Lett.* 2020;102:106096.

21. Hu YT, Bondarenko NP, Shieh CT, Yang CF. Traces and inverse nodal problems for Dirac-type integro-differential operators on a graph. *Appl Math Comput*. 2019;363:124606.
22. Koyunbakan H. The inverse nodal problem for a differential operator with an eigenvalue in the boundary condition. *Appl Math Lett*. 2008;21(12):1301-1305.
23. Guo Y, Wei G. Inverse problems: Dense nodal subset on an interior subinterval. *J Differ Equ*. 2013;255(7):2002-2017.
24. Albeverio S, Hryniv RO, Nizhnik LP. Inverse spectral problems for non-local Sturm–Liouville operators. *Inverse Probl*. 2007;23(2):523.
25. Nizhnik L. Inverse nonlocal Sturm–Liouville problem. *Inverse Probl*. 2010;26(12):125006.
26. Xu XJ, Yang CF. Inverse nodal problem for nonlocal differential operators. *Tamkang J Math*. 2019;50(3):337-347.
27. Hu YT, Yang CF, Xu XC. Inverse nodal problems for the Sturm–Liouville operator with nonlocal integral conditions. *J Inverse Ill-Posed Probl*. 2017;25(6):799-806.
28. Qin X, Gao Y, Yang C. Inverse Nodal Problems for the Sturm-Liouville Operator with Some Nonlocal Integral Conditions. *J Appl Math Phys*. 2019;7(01):111.
29. Yang CF. Inverse nodal problem for a class of nonlocal Sturm-Liouville operator. *Math Model Anal*. 2010;15(3):383-392.
30. Keskin B, Ozkan AS. Inverse nodal problems for Dirac-type integro-differential operators. *J Differ Equ*. 2017;263(12):8838-8847.
31. Bondarenko N, Buterin S. On recovering the Dirac operator with an integral delay from the spectrum. *RM*. 2017;71(3):1521-1529.
32. Gulsen T, Yilmaz E, Koyunbakan H. Inverse nodal problem for p-laplacian dirac system. *Math Methods Appl Sci*. 2017;40(7):2329-2335.
33. Guo Y, Wei G. Inverse nodal problem for Dirac equations with boundary conditions polynomially dependent on the spectral parameter. *RM*. 2015;67(1):95-110.
34. Wang YP, Yurko VA, Shieh CT. The uniqueness in inverse problems for Dirac operators with the interior twin-dense nodal subset. *J Math Anal Appl*. 2019;479(1):1383-1402.
35. Yang CF, Pivovarchik VN. Inverse nodal problem for Dirac system with spectral parameter in boundary conditions. *Compl Anal Oper Theory*. 2013;7(4):1211-1230.
36. Yang CF, Huang ZY. Reconstruction of the Dirac operator from nodal data. *Integr Equ Oper Theory*. 2010;66(4):539-551.
37. Abdullaev TS, Nabiev IMO. An algorithm for reconstructing the Dirac operator with a spectral parameter in the boundary condition. *Comput Math Math Phys*. 2016;56(2):256-262.
38. Ferzullazadeh AG, Nabiev IM. Some properties of the spectrum of the Dirac operator with a spectral parameter in the boundary condition. *Proc of Institute of Math and Mech of NAS of Azerbaijan*. 2020;46(2):189-196.
39. Ferzullazadeh AG. Solution algorithm of the inverse spectral problem for Dirac operator with a spectral parameter in the boundary condition. *Oper Matrices*. 2022;16(1):113-122.
40. Yang CF, Yurko V. Recovering Dirac operator with nonlocal boundary conditions. *J Math Anal Appl*. 2016;440(1):155-166.

**How to cite this article:** Ozkan AS, Adalar İ. Inverse nodal problem for Dirac operator with integral type nonlocal boundary conditions. *Math Meth Appl Sci*. 2023;46(1):986-993. doi:10.1002/mma.8561