

Trichoderma atroviride suppresses *Fusarium graminearum* by altering primary and secondary metabolite biosynthesis profiling

Evrım Özkale¹  | Emre Yörük²  | Mahir Budak³ | Ertan Mahir Korkmaz³ 

¹Department of Biology, Faculty of Arts and Sciences, Manisa Celal Bayar University, Manisa, Turkey

²Department of Molecular Biology and Genetics, Faculty of Arts and Sciences, Istanbul Yeni Yuzyil University, Istanbul, Turkey

³Department of Molecular Biology and Genetics, Faculty of Sciences, Sivas Cumhuriyet University, Sivas, Turkey

Correspondence

Evrım Özkale, Department of Biology, Faculty of Arts and Sciences, Manisa Celal Bayar University, Manisa, Turkey.
Email: evrimmiko@gmail.com

Funding information

Manisa Celal Bayar University Scientific Research Commission, Grant/Award Number: 2021-061

Abstract

The use of *Trichoderma* spp. offers an ecologically friendly tool for the struggle with mycotoxigenic *Fusarium* spp. Here, the alterations in transcriptome level were investigated in 6-day-old *Fusarium graminearum* cultures treated with *Trichoderma atroviride* and nontreated with *T. atroviride* (FGc), using whole transcriptome sequencing to better understand the associated biological processes. Transcriptome analysis indicated a total of 55 up-regulated genes and 728 down-regulated genes with $p_{adj} < 0.05$. Enrichment analysis revealed that the up-regulated genes were related to fatty acid biosynthesis, AMP-dependent biosynthesis, amino acid recognition/activation processes and secondary metabolite production, whereas down-regulated genes were involved in amino acid synthesis, oxidation–reduction processes, metal ion-binding and metabolic/catalytic activities. Among the down-regulated genes, the expression of pigmentation-related genes such as *aurO*, *gip1* and *aurR2* was remarkable. Similarly, the expression levels of key enzyme-coding genes involved in deoxynivalenol mycotoxin production were significantly decreased in the range of -1.77 and -2.94 . For up-regulated genes, nonribosomal peptide synthetase and polyketide synthase genes were notably distinguished from the remaining down-regulated genes as these genes can be involved in biosynthesis of common secondary metabolites. The results clearly emphasize that *T. atroviride* repressed the biosynthesis of primary metabolites in *F. graminearum* while simultaneously up-regulating the expression of genes involved in the synthesis of secondary metabolites. This is the first report showing how *T. atroviride* leads to transcriptome alterations and the findings suggest that *T. atroviride* could serve as an effective fungus by employing a wide variety of strategies against phytopathogenic fungi.

KEYWORDS

disease management, *Fusarium graminearum*, Fusarium head blight, RNA-Seq analysis, *Trichoderma atroviride*

1 | INTRODUCTION

Fusarium graminearum, teleomorph *Gibberella zeae*, is the predominant causal agent of Fusarium head blight (FHB) disease of small grain cereals in many agro-ecological regions. This cosmopolitan

phytopathogenic fungus has led to epidemics that have cost billions of dollars due to contamination with mycotoxins such as deoxynivalenol and zearalenone as well as reductions in crop quality and quantity (Goswami & Kistler, 2004; Leplat et al., 2013; Lori et al., 2009; Matny, 2015; Miedaner et al., 2008). Its high level of phenotypic

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2023 The Authors. *Plant Pathology* published by John Wiley & Sons Ltd on behalf of British Society for Plant Pathology.

and genomic plasticity has led to investigations related to *F. graminearum* becoming an important area of research in phytopathology. Moreover, the completed *F. graminearum* genome project has accelerated studies linked to FHB management (Cuomo et al., 2007; Pasquali & Migheli, 2014; Waalwijk et al., 2017; Wang et al., 2011).

F. graminearum is a soilborne phytopathogenic fungus that can persist in the soil for a long time, with a wide distribution pattern worldwide. The predominant strategy for managing *F. graminearum* involves the use of synthetic chemical pesticides. Even if common fungicides provide efficient management of FHB in a short time, there are several disadvantages, such as the development of fungicide resistance and deleterious environmental effects (Alabouvette et al., 2009; Chung et al., 2008; Özsoy et al., 2020; Zheng et al., 2015). In this context, effective and ecologically friendly management strategies, for example the use of plant-derived secondary metabolites such as terpenes, phenylpropanoids, polyketides and alkaloids, are required.

Microbial biological control agents (BCAs) have been frequently used in the fight against *Fusarium* diseases in recent years (Franco et al., 2011; Nourozian et al., 2006). Several fungal species such as *Gliocladium roseum* and *G. catenulatum* may be useful in the in vitro and in vivo control of *F. graminearum* diseases (Er, 2020; Legrand et al., 2018). *Trichoderma* spp. are notable among filamentous fungi for their ability to adapt to a wide variety of ecological conditions and lifestyles, as well as their benefits to plants (Zeilinger et al., 2016). *Trichoderma* species are also important mycoparasitic fungi and compete with other microorganisms for nutrients. *Trichoderma* species including *T. afroharzianum*, *T. atroviride*, *T. harzianum*, *T. virens* and *T. viride* have been used to combat *Fusarium* species. However, the majority of these studies related to interactions between *Trichoderma* spp. and *Fusarium* spp. are limited to two major fungi: *T. harzianum* and *F. oxysporum* (Amin et al., 2010; Mohamed & Haggag, 2006; Rawat et al., 2013; Strauch et al., 2011). Antagonistic effects of *Trichoderma* spp. on *F. graminearum* are only presented in a limited number of investigations. Among these, *T. atroviride* could be evaluated as a potential antagonistic agent against *F. graminearum*, as evidenced by reduced fungal biomass, decreased macroconidium quantity, induced oxidative stress and apoptosis-like-process (Błaszczuk et al., 2017; Cabrera et al., 2020; Matarese et al., 2012; Yörük et al., 2022). However, the response of *F. graminearum* treated with *T. atroviride* at the transcriptomic and/or metabolomic level is not available in the literature. Here, we aimed to reveal *T. atroviride*-induced transcriptomic alterations in *F. graminearum* and to elucidate the biological processes underlying these changes.

2 | MATERIALS AND METHODS

2.1 | In vitro fungal growth

F. graminearum PH-1 and *T. atroviride* TR8 strains were provided by Dr Tapani Yli-Mattila (Finland) and Dr Evrim Özkale (Turkey), respectively; the former was used as the phytopathogen and the latter as the BCA. Fungal cultures were conducted on potato dextrose agar (PDA) at $26 \pm 2^\circ\text{C}$ for 6 days. Plugs of mycelium (0.25 cm^2) of *F. graminearum*

PH-1 and *T. atroviride* TR8 were transferred onto the middle of the PDA plates. The control set (FGc) included only *F. graminearum* PH-1 grown on PDA, while the experiment set (FGvsTA) consisted of two Petri dishes including both *F. graminearum* PH-1 and *T. atroviride* TR8 cultures. Experimental cultures were conducted as sandwich cultures, in which the Petri dishes (without lids) belonging to two different species were brought together and sealed by Parafilm to form a single plate. Linear growth rates (LGR) of FGc and FGvsTA were measured as mm/day at the end of the seventh day of incubation (Yörük et al., 2022).

2.2 | Total RNA isolation, cDNA-second-strand cDNA synthesis, library preparation and next-generation sequencing

Total RNA was isolated from 7-day-old cultures using a RNeasy Plant Mini Kit (QIAGEN). Fresh mycelial material (50mg) was first ground with liquid nitrogen using a sterile pestle and mortar. After the homogenization process, total RNA was obtained following the manufacturer's recommendations. Genomic DNA was eliminated by DNase I (QIAGEN) treatment. The quantity and quality of total RNA were checked by Qubit RNA HS Assay Kit (Thermo) and agarose gel (0.8%). RNA integrity was evaluated by Bioptic Qsep100 (GC Biotech). Total RNA samples with sufficient quality were used to prepare cDNA libraries, in equal amounts per sample. Three libraries (one for the control set and two replicates for the experimental culture) were constructed for transcriptome sequencing.

The total RNA was used for first-strand cDNA synthesis with the poly(A) primer using a commercial first-strand cDNA synthesis kit (NEB) including M-MuLV enzyme according to the manufacturer's instructions. After quality check on agarose gel and Qubit fluorometer (Invitrogen-Thermo), the first-strand cDNA was transformed into double-strand (ds) cDNA by using RNase H and DNA polymerase I. Subsequently, the cDNA library fragments of desirable lengths were purified with the AMPure XP beads (Beckman Coulter). The cDNA libraries were then prepared for Illumina sequencing pair-end libraries with insert size of 300bp using the Illumina DNA Preparation Kit (Illumina Inc.). Thereafter, they were checked and normalized to 10nM by using Qubit 4.0 and agarose gel (0.8%) electrophoresis. Sequencing was performed on the MiSeq platform (Illumina), conducted at Advanced Technology Application and Research Centre of Sivas Cumhuriyet University (CUTAM, Turkey). The raw data were stored in the NCBI Sequence Read Archive (SRA) under accession number PRJNA970182.

2.3 | Transcriptome data analysis

All sequenced libraries were evaluated in terms of their quality and quantity by using Linux-based tools. The supercomputers with Linux system from TRUBA (Turkish Science e-Infrastructure, TUBITAK-Turkey) sources were used in next-generation sequencing analysis. Sequencing libraries obtained were first subjected to FastQC (and MultiQC) assays in order to check the quality of the reads (<http://>

www.bioinformatics.babraham.ac.uk/projects/fastqc/). The trimming process of raw reads was carried out using fastp preprocessor (Chen et al., 2018). After sequencing data were trimmed, the sequenced libraries were used in reference mapping by HiSat2 (Kim et al., 2015). The reads were then mapped to coding regions and counted by using HTseq, Kallisto and StringTie software (Anders et al., 2015; Bray et al., 2016; Pertea et al., 2015). The count tables were transferred to further assays.

2.4 | R-based differentially expressed gene analysis

The count tables belonging to next-generation sequencing (NGS) reads were used in data mining processes by R software v. 4.2.2. DESeq2, apegm, pheatmap, tidyverse, ggrepel and ashR packages were installed by CRAN or Bioconductor. Count files from previous analyses were normalized using the DESeq2 package. Common commands were used in order to obtain the volcano plot, heatmaps and differentially expressed gene (DEG) table. The genes with an adjusted p value (p_{adj}) ≤ 0.05 and a \log_2 fold change ≥ 1 were accepted/considered as differentially expressed genes. Genes with significantly altered expressions were transferred to functional analysis.

2.5 | Gene ontology and KEGG orthology analysis

Unigenes were analysed with BLASTX algorithm using NCBI, Swiss-Prot (http://web.expasy.org/docs/swissprot_guideline.html),

Kyoto Encyclopedia of Genes and Genomes (KEGG) and Cluster of Orthologous Groups (COG) (<http://www.ncbi.nlm.nih.gov/COG/>) databases. Protein functions were annotated and coding sequence (CDS) regions were confirmed by ETSScan. Unigenes were also subjected to Blast2GO assays. The functional annotations and classifications were carried out using Gene Ontology (GO) (<http://geneontology.org/>) and WEGO software (<https://wego.genomics.cn/>). GO trees were constructed from GO Biological process pathway databases with the parameters of false discovery rate (FDR) cut-off value as 0.05 and aspect ratio as 2.0. Blastall was used in obtaining the functional annotations of unigenes by COG and KEGG databases. Metabolic pathway annotations were obtained by KEGG (<http://www.genome.jp/kegg/pathway.html>). Enrichment analyses were also carried out by GoShiny online tool (<http://bioinformatics.sdstate.edu/go>).

3 | RESULTS

3.1 | In vitro fungal growth, total RNA isolation, cDNA and second-strand cDNA synthesis

Linear growth rate (LGR) of FGvsTA was compared with FGc. The decreased in vitro growth capacity was obtained by the end of the seventh day of incubation. LGR values of FGc and FGvsTA sets were 12.86 ± 0.01 and 7.71 ± 1.82 mm/day, respectively. A significant decrease in LGR was detected in the FGvsTA set ($***p < 0.01$). Total RNA was isolated from control and experiment sets at a quantity




FIGURE 1 Volcano plot generated based on \log_2 fold change in *Fusarium graminearum* cultures treated with *Trichoderma atroviride*.

of 0.5–2 µg/µL and absorbance ratio values of 1.9–2.0 $A_{260/280}$. The quality and the quantity of cDNA and ds cDNA molecules were in the ranges as reported by Nagalakshmi et al. (2010).

3.2 | Transcriptome-wide analysis in *F. graminearum*

Whole transcriptome data for *F. graminearum* PH-1 treated or non-treated with *T. atroviride* were generated to reveal the transcriptomic alterations in the phytopathogenic fungus under the biotic stress factor. Linux-based data were completed by obtaining the count table including Gene-ID numbers, \log_2 fold changes with *SE* values, *p* values and p_{adj} values as a single .tsv file (File S1). $p_{adj} \leq 0.05$ was accepted as reference value for identifying the statistically significant differences in genes with differential expression. Minimum p_{adj} values were recorded as $6.8e-21$ (FGSG_04745) for down-regulated genes and $4.6e-08$ for up-regulated genes (FGSG_11455). In total, 55 up-regulated genes with $p_{adj} \leq 0.05$ were detected, whereas 728 genes exhibited a significant decrease between FGc and FGvsTA sets (File S1). A volcano plot indicated that the majority of the transcripts were not significantly altered (Figure 1). Table 1 highlights the top 10

up-regulated and down-regulated genes with the highest \log_2 fold change (FC) values. Up-regulated genes with dramatic fold changes are found in a wide range of different pathways such as actinorhodin (polyketide antibiotic) biosynthetic pathway (FGSG_11455), shikimate pathway (FGSG_11457) and citric acid cycle (FGSG_10444). Table 1 also shows similar striking findings for up-regulated genes involved in antifungal protein synthesis (FGSG_04745), oxidative metabolism of hydrocarbons (FGSG_00043) and sugar transporters (FGSG_05882). DEG analysis revealed that *T. atroviride* led to significant alterations in expression of genes involved in the secondary metabolite biosynthesis in *F. graminearum*. The overall major transcriptomic changes between FGc and FGvsTA and the dramatically up-regulated and/or down-regulated genes are shown in Figures 2 and 3, respectively. Two categorically different GO trees were obtained by Gene Ontology analysis (Figure 4). The GO tree belonging to up-regulated genes included genes predominantly linked to secondary metabolite production processes and fatty acid synthesis (Figure 4a), while the GO tree of down-regulated genes harboured the genes involved in primary metabolite biosynthesis, oxidation-reduction processes and metabolic/catalytic activities (Figure 4b). In this way, the GO trees facilitated the differentiation of different biological processes that involve genes with different critical levels

TABLE 1 Top 20 altered genes in *Fusarium graminearum* treated with *Trichoderma atroviride*.

Gene ID	\log_2 FC	p_{adj}	Annotation
Top 10 up-regulated genes			
FGSG_11455	4.96	$4.6e-08$	Dimeric α - β barrel (IPR011008)
FGSG_10443	4.16	$5.2e-08$	Hypothetical protein
FGSG_10442	3.14	$1.0e-05$	Hypothetical protein
FGSG_11457	3.05	$8.3e-05$	Mmf1/domain is similar in structure to chorismate mutase (IPR006175)
FGSG_10444	2.38	$3.0e-04$	L-lactate/malate dehydrogenase (PIRSF000102)
FGSG_05048	2.35	$4.6e-04$	Dihydrodipicolinate synthase (PIRSF001365)
FGSG_13521	2.31	$4.0e-04$	Hypothetical protein
FGSG_10445	2.17	$1.5e-03$	Phosphoenolpyruvate synthase (IPR006319)
FGSG_10922	2.07	$1.5e-03$	Hypothetical protein (with expansin domains [IPR007117, IPR007112])
FGSG_12977	1.94	$4.1e-03$	Nucleoside triphosphate hydrolase (IPR027417)
Top 10 down-regulated genes			
FGSG_04745	-6.29	$6.8e-21$	Antifungal protein domain superfamily (IPR023112)
FGSG_09066	-4.68	$7.9e-13$	Hypothetical protein (with signal peptide)
FGSG_00043	-4.05	$2.5e-09$	FAD/NAD(P)-binding domain superfamily (IPR036188)
FGSG_07988	-3.58	$4.6e-08$	Hypothetical protein (with signal peptide)
FGSG_13979	-3.81	$5.3e-08$	Acyl-CoA <i>N</i> -acyltransferase (IPR016181)
FGSG_07896	-3.61	$1.6e-08$	Chloramphenicol acetyltransferase-like domain superfamily (IPR023213)
FGSG_04527	-3.61	$3.0e-08$	Carboxypeptidase S1 (IPR001563)
FGSG_08083	-3.45	$3.1e-08$	Pyridoxal phosphate-dependent decarboxylase (IPR002129)
FGSG_05882	-3.54	$4.6e-08$	Major facilitator, sugar transporter-like (IPR005828)
FGSG_00053	-3.50	$3.5e-08$	Aryl-alcohol dehydrogenase (IPR023210)

of expression. Enrichment assays revealed that up-regulated genes were primarily related to secondary metabolite production steps, whereas down-regulated genes were related to main metabolite synthesis and biological processes with great impact such as amino acid transport, ion transport and oxidation–reduction processes (Figure 5). Table 2 shows the specific genes with significantly altered expression according to enrichment analysis. The enrichment analysis equipped with InterPro database and/or AmiGO was collected as a single.xlsx file consisting of the list of up-regulated and down-regulated genes (File S2).

3.3 | Alterations in expression of genes related to prevalent secondary metabolite biosynthesis

The expression levels of genes encoding key enzymes that regulate biosynthesis of common secondary metabolites highlight the presence of significant changes in *F. graminearum* (Table 3). Moreover, the results of DEG analysis were also considered for the genes related to DON synthesis, aurofusarin synthesis, zearalenone synthesis, apoptosis and autophagy (Figure 6). Figure 6a and File S1 show that majority of the genes involved in apoptosis-like processes did not alter as a result of *T. atroviride* treatment. Similarly, autophagy-related genes did not significantly alter in the FGvsTA set (Figure 6b). However, pigmentation-related aurofusarin biosynthesis genes were significantly down-regulated in response to *T. atroviride* (Figure 6c). Similarly, the genes located in the core synthetic cluster of DON production were significantly decreased (Figure 6d). There were no co-alterations in zearalenone biosynthesis-related genes (Figure 6e). While butanolide production-associated CYP67 showed down-regulation, up-regulation was observed in some other related key genes of secondary metabolite production (Figure 6f).

4 | DISCUSSION

Today, the whole genome of any kind of organism including large/giant genomes such as *Homo sapiens* could be sequenced in a single day. NGS strategies have evolved from the first-generation whole genome sequencing strategies to third- or fourth-generation sequencing strategies such as Nanopore Sequencing or VASA-Seq transcriptome analysis within the last two decades (Behjati & Tarpey, 2013; Branton et al., 2008; Salmen et al., 2022). RNA-Seq technology uses large-scale parallel sequencing of short cDNA molecules converted from RNA and provides transcriptome reads with high resolution within a short time in comparison with the classical Sanger dideoxy termination method and microarray-based sequencing technologies. Qualitative and quantitative analyses including exon-intron boundaries, haplotyping and transcript levels can be easily carried out by mapping reads to reference genome files (Nagalakshmi et al., 2010). Here, cDNA-based transcriptome analysis in *F. graminearum* treated with *T. atroviride* was carried out in order to discover which strategies the phytopathogen might use in response to a biotic stress factor.

Many of the studies in which *F. graminearum* was the subject of research have been focused on chemotyping, genetic diversity and determination of common fungicide resistance level (Miedaner et al., 2008; Pasquali & Migheli, 2014; Przemieniecki et al., 2014; Wang et al., 2011; Yang et al., 2018; Yörük et al., 2018; Yörük & Yli-Mattila, 2019). Studies on *F. graminearum* at the transcript level have been restricted to alterations of expression of the genes located on the *tri5* gene cluster in response to abiotic stress factors (Boutigny et al., 2009; Gazdağlı et al., 2018; Jiao et al., 2008; Kulik et al., 2014; Marín et al., 2010; Merhej et al., 2010; Oufensou et al., 2020; Pinson-Gadais et al., 2008; Yörük, 2018). There are a few investigations related to transcriptome-wide analysis in *F. graminearum*. Affymetrix




FIGURE 2 Major transcriptomic changes related to primary or secondary biological processes in *Fusarium graminearum* cultures treated with *Trichoderma atroviride* (FGvsTA) compared with untreated *Fusarium graminearum* (FGc). [Colour figure can be viewed at wileyonlinelibrary.com]




FIGURE 3 Dramatically up-regulated (a) and down-regulated (b) genes. FGc1, FGc2: untreated *Fusarium graminearum* control sets 1 and 2. FGvsTA1, FGvsTA2: *Fusarium graminearum* cultures treated with *Trichoderma atroviride*, sets 1 and 2. [Colour figure can be viewed at [wileyonlinelibrary.com](https://onlinelibrary.wiley.com/doi/10.1111/ppa.13768)]





FIGURE 4 Gene Ontology (GO) trees related to up-regulated (a) and down-regulated (b) genes in *Fusarium graminearum* cultures treated with *Trichoderma atroviride*. [Colour figure can be viewed at wileyonlinelibrary.com]

chips and DeepSAGE strategies were developed and conducted as suitable tools for both in vitro and in planta studies in *F. graminearum* (Güldener et al., 2006; Liu et al., 2010). However, these approaches were not followed after high-throughput RNA-Seq-based strategies became more common. Zhao et al. (2013) studied gene annotations in *F. graminearum* by RNA-Seq strategy and reported that alternative splicing strategy processes are regulated by development phases. A holistic approach equipped with RNA-Seq was used in order to reveal the alterations in individual genes or genes located on gene clusters related to secondary metabolite biosynthesis (Sieber et al., 2014). This study revealed that a total of 87 different gene clusters involved in the biosynthesis of known or unknown secondary metabolites showed notable and strong alterations in gene expression specific to enzymatic activity (File S1). Zapparata et al. (2021) investigated

the transcriptomic changes in *Trichoderma gamsii* and *F. graminearum* in dual confrontation cultures at the sensing phase of growth. They reported that *F. graminearum* up-regulates the expression of genes related to transmembrane transporters and killer toxins. In total, 653 up-regulated and 75 down-regulated genes with significantly altered expression ($p_{adj} \leq 0.05$ and \log_2FC for >1) were recorded in *F. graminearum* treated with *T. gamsii*. Interestingly, they reported both down-regulation and up-regulation in genes coding for metabolites or ion transporter enzymes, with dramatic up-regulation for MFS general substrate transporter (FGRA07_11673), sugar transporter (FGRA07_06312) and auxin efflux carrier (FGRA07_02452), and notable down-regulation for sugar transporter (FGRA07_11651) and sulphur transport protein (FGRA07_10600) in *F. graminearum* treated with *T. gamsii*.

Up-regulated genes in Enrichment Analysis



Down-regulated genes in Enrichment Analysis




FIGURE 5 Specific transcripts related to characteristic pathways obtained from enrichment analysis. [Colour figure can be viewed at wileyonlinelibrary.com]

Dual culture tests including *F. graminearum* against *T. gamsii* concluded at the sensing phase of fungi (Zapparata et al., 2021). Unlike the mycoparasitism strategy (dual confrontation cultures, "before contact") used by Zapparata et al. (2021), sandwich cultures, which have been preferred in previous studies (Ramangouda et al., 2023; Yörük et al., 2022), were used in this study. These cultures can suppress in vitro growth of *F. graminearum* by up to 40%–50% for 7 days. Zapparata et al. (2021) used *T. gamsii* as BCA, which could be accepted as a strong BCA against *Fusarium* spp. (Baroncelli et al., 2016; Matarese et al., 2012). Here, *T. atroviride* was selected because it has recently been used as a BCA against *Fusarium* spp. (Cabrera et al., 2020; Coninck et al., 2020; Yörük et al., 2022). To

our knowledge, this is the first report to show the transcriptome-wide alterations in *F. graminearum* treated with *T. atroviride* under sandwich cultures that could induce the volatile organic compound-based antagonistic effects of *Trichoderma* spp. Here, the number of down-regulated genes in *F. graminearum* was found to be higher than that of a recent report by Zapparata et al. (2021). This may have resulted from the BCA species used or may be due to the mycoparasitism strategy chosen. However, similar transcript-changing profiles were obtained in terms of down-regulation of genes coding for ion and/or metabolite transporter enzymes.

The data obtained from this study contribute novel information to the field by providing a unique perspective on transcript

TABLE 2 Specific genes with statistically significant alterations obtained from enrichment analysis in *Fusarium graminearum* PH-1 treated with *Trichoderma atroviride*.

Gene-ID	Gene	Log ₂ FC
Up-regulated genes		
FGSG_02394	NPS15 (nonribosomal peptide synthetase)	1.42
FGSG_11026	NPS1 (nonribosomal peptide synthetase)	1.66
FGSG_11395	NPS14 (nonribosomal peptide synthetase)	1.24
FGSG_11659	NPS8/GRA1 (nonribosomal peptide synthetase)	1.02
FGSG_13783	NPS18 (nonribosomal peptide synthetase)	1.09
FGSG_13878	FgNRPS5 (nonribosomal peptide synthetase)	1.24
FGSG_01790	PKS11 (polyketide synthase)	1.01
FGSG_03340	PKS8 (polyketide synthase)	1.08
FGSG_03964	PKS14 (polyketide synthase)	1.05
FGSG_04694	PKS2 (polyketide synthase)	1.28
FGSG_09182	PKS3 (polyketide synthase)	1.46
FGSG_12126	ZEA2/PKS13 (reducing polyketide synthase)	1.31
FGSG_06645	FGSG_06645 (hypothetical protein)	1.14
Down-regulated genes		
FGSG_02321	aurO (aurofusarin biosynthesis)	-3.09
FGSG_02327	aurF (aurofusarin biosynthesis)	-2.92
FGSG_02328	gip1 (aurofusarin biosynthesis)	-2.82
FGSG_02329	Hypothetical protein with FAS domain	-3.02
FGSG_02323	aurR2 (aurofusarin biosynthesis)	-2.04
FGSG_02326	aurJ (aurofusarin biosynthesis)	-2.81
FGSG_05491	Histone H4	-1.35
FGSG_00702	Nucleotide-diphospho-sugar transferase	-2.75
FGSG_00620	Ammonium transporter MEP2	-2.22
FGSG_01675	Dicarboxylic amino acid permease	-3.21
FGSG_02094	Ammonium/urea transporter	-2.71
FGSG_03412	Amino acid permease inda1	-2.15
FGSG_05277	Uracil permease	-1.94
FGSG_03028	RNA polymerase I-specific transcription initiation factor RRN9	-2.34
FGSG_06118	Related to integral membrane protein	-1.98
FGSG_07988	phiA (cell wall protein)	-3.58
FGSG_08122	afpA (alkaline foam protein A)	-1.63
FGSG_02320	aurR1 (aurofusarin biosynthesis)	-2.12
FGSG_05159	Iron transport multicopper oxidase FET3 precursor	-2.43

TABLE 2 (Continued)

Gene-ID	Gene	Log ₂ FC
FGSG_08079	Benzoate 4-monooxygenase	-3.09
FGSG_03412	Amino acid permease inda1 (03412)	-2.15
FGSG_04637	Amino acid permease inda1 (04637)	-2.90
FGSG_05574	gap1 (general amino acid permease)	-2.26
FGSG_09053	Amino acid permease (09053)	-2.56
FGSG_09297	Amino acid permease/SLC12A domain (09297)	-2.71
FGSG_00043	Putative thioredoxin reductase	-4.05
FGSG_08077	NADPH dehydrogenase YqjM-like	-1.80
FGSG_08079	Cytochrome P450, conserved site	-3.09
FGSG_08081	Isopenicillin N synthase-like, Fe ²⁺ 2OG dioxygenase domain (2og-fe)	-3.33
FGSG_02322	aurT (aurofusarin biosynthesis)	-1.92
FGSG_05882	MFS transporter superfamily	-3.54
FGSG_09374	Acetate transporter GPR1/FUN34/SatP family	-1.84
FGSG_03319	Cell division cycle ATPase	-2.09
FGSG_08083	Pyridoxal phosphate-dependent transferase	-3.45
FGSG_00049	Branched-chain amino acid aminotransferase II	-2.49
FGSG_08082	Hypothetical protein with acyl-CoA N-acyltransferase	-2.75
FGSG_08852	Putative cryptochrome DASH	-3.20
FGSG_09690	Peptidyl-prolyl isomerase	-2.54
FGSG_05160	Iron permease FTR1	-2.66
FGSG_08721	SODC_NEUCR superoxide dismutase	-2.23
FGSG_02523	Mss4/translationally controlled tumour-associated protein TCTP	-2.16
FGSG_02111	Cytochrome P450 (02111)	-2.30
FGSG_02117	Cytochrome P450 (02117)	-2.73
FGSG_02273	Aldehyde dehydrogenase	-2.56
FGSG_03191	Cytochrome P450 (3191)	-2.42
FGSG_03535	tri4 (trichodiene oxygenase)	-2.95
FGSG_03569	gaoB (galactose oxidase precursor)	-1.46
FGSG_04126	Uricase	-2.30
FGSG_05322	Fatty acid synthase subunit β dehydratase	-1.35
FGSG_09879	Cytochrome c oxidase, subunit VIIa	-1.88
FGSG_10677	Copper amine oxidase	-2.13

profiling of genes involved in secondary metabolite biosynthesis. Genes coding for polyketide synthase were found as mostly up-regulated in response to the presence of biotic stress (Figure 5a). Overall, at least 15 nonribosomal peptide synthase and polyketide synthase genes were up-regulated, representing a sign of the *F. graminearum* survival strategy (Table 2). Among these genes, *nmps* and *pks* genes could be associated with more than one

TABLE 3 Alterations in the genes involved in common secondary metabolite biosynthesis in *Fusarium graminearum* PH-1 treated with *Trichoderma atroviride*.

Metabolite	Gene ID	Gene	Role	Fold change
Butenolide	FGSG_08079	CYP67	CYP67-like; cytochrome P450 family 67 and similar cytochrome P450s	-3.09
Malonichrome	FGSG_11026	NPS1	Extracellular siderophore, induced in planta	1.66
Ferricrocin	FGSG_05372	NPS2	Intracellular siderophore	1.03
Zearalenone	FGSG_02395 FGSG_12055	PKS13	Polyketide synthase; powerful xenoestrogen in animals, no effect on virulence	1.08
Fusarielin	FGSG_10464	PKS9	Polyketide synthase	1.56
Aurofusarin	FGSG_02320	aurR1	Golden yellow/red pigment of mycelium, low toxicity (high concentrations in feed can affect antioxidant levels in eggs)	-2.12
Orcinol	FGSG_03964	PKS28	Responsible for production of orsellinic acid/orcinol	1.05
Deoxynivalenol	FGSG_03537	TRI5	Protein biosynthesis inhibitor, virulence factor on wheat	-1.78

secondary metabolite production in phytopathogenic fungi (Frandsen et al., 2006; Gaffoor et al., 2005; Gazdađlı et al., 2023; Ma et al., 2013). Even if the majority of these *nrps* and *pks* genes were associated with the precise pathways, up-regulated genes (such as FGSG_11026, FGSG_02394, FGSG_09182 and FGSG_04694; Table 2) in the FGvsTA experimental set should be used in further studies in evaluating the pathogenomics and pathotranscriptomics in *Fusarium*. Essential genes (*tri4*, *tri5*, *tri9* and *tri14*) related to DON production were down-regulated in the FGvsTA set (Figure 5). The genes of *tri4*, a multifunctional oxygenase responsible for catalysis of four steps in DON synthesis and *tri5*, trichodiene synthase, responsible for the first step in DON biosynthesis, could be regarded as marker genes (Yörük & Yli-Mattila, 2019). The majority of previous studies on the potential decrease in DON biosynthesis focused on the identification of changes in the *tri4* and *tri5* genes (Boutigny et al., 2009, 2010; Merhej et al., 2011; Pinson-Gadais et al., 2008; Ponts et al., 2007; Yaguchi et al., 2009; Yörük & Albayrak, 2019; Yörük et al., 2022). In this manner, a significant decrease in *tri4* and *tri5* genes in FGvsTA experiment sets yielded important data in terms of revealing that *T. atroviride* could be used as a potential effective BCA against *F. graminearum* and related species in further studies. In addition to gene expression alterations related to DON biosynthesis, butanolide mycotoxin biosynthesis-related CYP67 was also found as down-regulated. However, total transcript changes were not detected in the genes involved in zearalenone

production except for *PKS13*. Among the down-regulated genes, the expression of genes linked to aurofusarin biosynthesis had notable expression changes (Figure 6). Almost all genes located on the *AUR* gene cluster were down-regulated. These transcriptomic changes were supported by the Petri dish profiling of the FGvsTA set (Figure 2). It is clear that the lifecycle of *F. graminearum* was disrupted by *T. atroviride* because aurofusarin and its precursors are primarily responsible for the pigmentation and antifungal effects of *F. graminearum* (Frandsen et al., 2006; Gaffoor et al., 2005; Gazdađlı et al., 2023). Additionally, killer toxin genes (FGSG_10551, FGSG_00061, FGSG_00060 and FGSG_00062) exhibited down-regulation or no expression in the FGvsTA set in contrast to the previous report including dual culture tests in *F. graminearum* (Zapparata et al., 2021).

As mentioned earlier, the enrichment analysis showed that many of the genes with up-regulation were associated with secondary metabolite biosynthesis processes. However, here, both GO and KEGG analyses revealed that the down-regulated genes (Figure 5b) belonged to distinct biological processes that are crucial for the survival of *F. graminearum*, such as amino acid metabolism (e.g., FGSG_09053 and FGSG_04637), ammonium/urea transport (e.g., FGSG_02094 and FGSG_00620), ion transport (e.g., FGSG_05159.1 and FGSG_08081) and oxidation-reduction processes (e.g., FGSG_00043 and FGSG_03569). The dramatic alterations (especially down-regulation) in these kinds of genes,




FIGURE 6 Alterations in genes related to apoptosis (a), autophagy (b), aurofusarin synthesis (c), deoxynivalenol synthesis (d), zearalenone synthesis (e) and key enzymes in secondary metabolites (f). Significant difference at * $p_{adj} < 0.05$, ** $p_{adj} < 0.01$ and *** $p_{adj} < 0.001$. [Colour figure can be viewed at wileyonlinelibrary.com]

which could be associated with primary metabolism, could be indicative of a poor quality of lifecycle in *F. graminearum*. However, it appears that about 5.3% of down-regulated genes (with p_{adj} value ≤ 0.05 and a \log_2 fold change > 1); 728 of a total of 13,694 genes) could not lead *F. graminearum* to programmed cell death (PCD) arising from apoptosis-like process and autophagy (Figure 6). Moreover, no cross-talk was present between apoptosis and autophagy according to the DEG analysis result. Apart from PCD related to apoptosis-like process and autophagy, necroptosis and/or pyroptosis can also initiate PCD in fungi (Gonçalves et al., 2017). However, there is still a lack of information about the genes and pathways involved in these alternate PCD strategies. Necrosis-like cell death regulatory proteins like NWD2 and PNT1 have yet to be annotated and characterized, despite the presence of genomes of *F. graminearum* PH-1 (GCA_000240135.3) and *F. graminearum* ITEM124 (GCA_002352725.1) in databases. We hypothesize that PCD via necroptosis or pyroptosis could be a possible strategy chosen by *F. graminearum* when it is exposed to biotic stress. Further annotation studies are required to fully understand the alternative PCD pathways.

In conclusion, the comprehensive and in-depth investigations here reveal the mechanisms by which *T. atroviride* competes with *F. graminearum*. The majority of the studies on the struggle between *Trichoderma* spp. and *Fusarium* spp. have been conducted on phenotypic and analytical tests (Baroncelli et al., 2016; Benítez et al., 2004; Matarese et al., 2012; Mohamed & Haggag, 2006; Sharma et al., 2013). In this study, transcriptomic alterations were investigated in *F. graminearum* treated with *T. atroviride* in sandwich cultures. It was shown that *T. atroviride* decreases the expression of genes involved in key biological pathways in *F. graminearum*. This was also demonstrated by a disruption in the abundance of transcripts related to the processes of mycotoxin biosynthesis, ion/metabolite transport, amino acid metabolism and oxidation-reduction. Based on the findings of this study, *T. atroviride* has the potential to be an effective BCA that employs a wide variety of strategies to fight against phytopathogenic microorganisms.

ACKNOWLEDGEMENTS

The authors are grateful to Dr Tapani Yli-Mattila from the University of Turku-Finland for providing *F. graminearum* PH-1 and to Dr Evrim Özkale from Manisa Celal Bayar University—Turkey for providing *T. atroviride* TR8. The members of Evolutionary Bioinformatics Research Groups (EBRG) from Sivas Cumhuriyet University are thanked for their contributions to laboratory work. We thank the anonymous reviewers for their careful reading of the manuscript and many constructive comments. This study was funded by the Scientific Research Commission of the Manisa Celal Bayar University (project 2021/061).

CONFLICT OF INTEREST STATEMENT

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

DATA AVAILABILITY STATEMENT

Raw sequence data that support this work have been deposited in NCBI BioProject at <https://www.ncbi.nlm.nih.gov/bioproject/> with accession number PRJNA970182.

ORCID

Evrin Özkale  <https://orcid.org/0000-0002-1707-9777>

Emre Yörük  <https://orcid.org/0000-0003-2770-0157>

Ertan Mahir Korkmaz  <https://orcid.org/0000-0003-0699-1354>

REFERENCES

- Alabouvette, C., Olivain, C., Migheli, Q. & Steinberg, C. (2009) Microbiological control of soil-borne phytopathogenic fungi with special emphasis on wilt-inducing *Fusarium oxysporum*. *New Phytologist*, 184, 529–544.
- Amin, F., Razdan, V.K., Mohiddin, F.A., Bhat, K.A. & Sheikh, P.A. (2010) Effect of volatile metabolites of *Trichoderma* species against seven fungal plant pathogens *in-vitro*. *Journal of Phytochemistry*, 2, 34–37.
- Anders, S., Pyl, P.T. & Huber, W. (2015) HTSeq—a python framework to work with high-throughput sequencing data. *Bioinformatics*, 31, 166–169.
- Baroncelli, R., Zapparata, A., Piaggieschi, G., Sarrocco, S. & Vannacci, G. (2016) Draft whole-genome sequence of *Trichoderma gamsii* T6085, a promising biocontrol agent of *Fusarium* head blight on wheat. *Genome Announcements*, 4, e01747-15.
- Behjati, S. & Tarpey, P.S. (2013) What is next generation sequencing? *Archives of Disease in Childhood – Education and Practice*, 98, 236–238.
- Benítez, T., Rincón, A.M., Limón, M.C. & Codon, A.C. (2004) Biocontrol mechanisms of *Trichoderma* strains. *International Microbiology*, 7, 249–260.
- Błaszczak, L., Basińska-Barczak, A., Ćwiek-Kupczyńska, H., Gromadzka, K., Popiel, D. & Stępień, Ł. (2017) Suppressive effect of *Trichoderma* spp. on toxigenic *Fusarium* species. *Polish Journal of Microbiology*, 66, 85–100.
- Boutigny, A.L., Atanasova-Pénichon, V., Benet, M., Barreau, C. & Richard-Forget, F. (2010) Natural phenolic acids from wheat bran inhibit *Fusarium culmorum* trichothecene biosynthesis *in vitro* by repressing *tri* gene expression. *European Journal of Plant Pathology*, 127, 275–286.
- Boutigny, A.L., Barreau, C., Atanasova-Pénichon, V., Verdal-Bonnin, M.N., Pinson-Gadais, L. & Richard-Forget, F. (2009) Ferulic acid, an efficient inhibitor of type B trichothecene biosynthesis and *tri* gene expression in *Fusarium* liquid cultures. *Mycological Research*, 113, 746–753.
- Branton, D., Deamer, D.W., Marziali, A., Bayley, H., Benner, S.A., Butler, T. et al. (2008) The potential and challenges of nanopore sequencing. *Nature Biotechnology*, 26, 1146–1153.
- Bray, N.L., Pimentel, H., Melsted, P. & Pachter, L. (2016) Near-optimal probabilistic RNA-seq quantification. *Nature Biotechnology*, 34, 525–527.
- Cabrera, M., Garmendia, G., Rufo, C., Pereyra, S. & Vero, S. (2020) *Trichoderma atroviride* as a biocontrol agent of *Fusarium* head blight by reducing the inoculum of the pathogen in wheat straw. *Terra Latinoamericana*, 38, 629–651.
- Chen, S., Zhou, Y., Chen, Y. & Gu, J. (2018) fastp: an ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics*, 34, i884–i890.
- Chung, W.H., Ishii, H.H., Nishimura, K., Ohshima, M., Iwama, T. & Yoshimatsu, H. (2008) Genetic analysis and PCR-based identification of major *Fusarium* species causing head blight on wheat in Japan. *Journal of General Plant Pathology*, 74, 364–374.
- Coninck, E., Scaufflaire, J., Gollier, M., Liénard, C., Foucart, G., Manssens, G. et al. (2020) *Trichoderma atroviride* as a promising biocontrol

- agent in seed coating for reducing *Fusarium* damping-off on maize. *Journal of Applied Microbiology*, 129, 637–651.
- Cuomo, C.A., Güldener, U., Xu, J.R., Trail, F., Turgeon, B.G., Di Pietro, A. et al. (2007) The *Fusarium graminearum* genome reveals a link between localized polymorphism and pathogen specialization. *Science*, 317, 1400–1402.
- Er, Y. (2020) Determination of antagonistic effect of bio-agents against cereal damping-off caused by *Fusarium graminearum* in wheat. *Egyptian Journal of Biological Pest Control*, 30, 43.
- Franco, T.S., Garcia, S., Hirooka, E.Y., Ono, Y.S. & Dos Santos, J.S. (2011) Lactic acid bacteria in the inhibition of *Fusarium graminearum* and deoxynivalenol detoxification. *Journal of Applied Microbiology*, 111, 739–748.
- Frandsen, R.J., Nielsen, N.J., Maolanon, N., Sørensen, J.C., Olsson, S., Nielsen, J. et al. (2006) The biosynthetic pathway for aurofusarin in *Fusarium graminearum* reveals a close link between the naphthoquinones and naphthopyrones. *Molecular Microbiology*, 61, 1069–1080.
- Gaffoor, I., Brown, D.W., Plattner, R., Proctor, R.H., Qi, W. & Trail, F. (2005) Functional analysis of the polyketide synthase genes in the filamentous fungus *Gibberella zeae* (anamorph *Fusarium graminearum*). *Eukaryotic Cell*, 4, 1926–1933.
- Gazdağlı, A., Baştumur, G.Y., Yörük, E., Özkorucuklu, S.P. & Albayrak, G. (2023) Molecular analysis of the genes responsible for catalysing intracellular steps of aurofusarin biosynthesis in *Fusarium culmorum*. *Brazilian Archives of Biology and Technology*, 66, e23210821.
- Gazdağlı, A., Sefer, Ö., Yörük, E., Varol, G.I., Teker, T. & Albayrak, G. (2018) Investigation of camphor effects on *Fusarium graminearum* and *F. culmorum* at different molecular levels. *Pathogens*, 7, 90–98.
- Gonçalves, A.P., Heller, J., Daskalov, A., Videira, A. & Glass, N.L. (2017) Regulated forms of cell death in fungi. *Frontiers in Microbiology*, 8, 1837.
- Goswami, R.S. & Kistler, H.C. (2004) Heading for disaster: *Fusarium graminearum* on cereal crops. *Molecular Plant Pathology*, 5, 515–525.
- Güldener, U., Seong, K.Y., Boddu, J., Cho, S., Trail, F., Xu, J.R. et al. (2006) Development of a *Fusarium graminearum* Affymetrix GeneChip for profiling fungal gene expression in vitro and in planta. *Fungal Genetics and Biology*, 43, 316–325.
- Jiao, F., Kawakami, A. & Nakajima, T. (2008) Effects of different carbon sources on trichothecene production and *tri* gene expression by *Fusarium graminearum* in liquid culture. *FEMS Microbiology Letters*, 285, 212–219.
- Kim, D., Langmead, B. & Salzberg, S.L. (2015) HISAT: a fast spliced aligner with low memory requirements. *Nature Methods*, 12, 357–360.
- Kulik, T., Buško, M., Pszczółkowska, A., Perkowski, J. & Okorski, A. (2014) Plant lignans inhibit growth and trichothecene biosynthesis in *Fusarium graminearum*. *Letters in Applied Microbiology*, 59, 99–107.
- Legrand, F., Picot, A., Cobo-Díaz, J.F. & Floch, G.L. (2018) Evaluation of the efficacy of a biocontrol agent, *Gliocladium catenulatum*, to colonise soils and to reduce *Fusarium graminearum* growth under microcosm and field conditions. *IOBC/WPRS Bulletin*, 135, 45–47.
- Leplat, J., Friberg, H., Abid, M. & Steinberg, C. (2013) Survival of *Fusarium graminearum*, the causal agent of Fusarium head blight: a review. *Agronomy for Sustainable Development*, 33, 97–111.
- Liu, X., Jiang, J., Shao, J., Yin, Y. & Ma, Z. (2010) Gene transcription profiling of *Fusarium graminearum* treated with an azole fungicide tebuconazole. *Applied Microbiology and Biotechnology*, 85, 1105–1114.
- Lori, G.A., Sisterna, M.N., Sarandón, S.J., Rizzo, I. & Chidichimo, H. (2009) Fusarium head blight in wheat: impact of tillage and other agronomic practices under natural infection. *Crop Protection*, 28, 495–502.
- Ma, L.J., Geiser, D.M., Proctor, R.H., Rooney, A.P., O'Donnell, K., Trail, F. et al. (2013) *Fusarium* pathogenomics. *Annual Review of Microbiology*, 67, 399–416.
- Marín, P., Jurado, M., Magan, N., Vázquez, C. & González-Jaén, M.T. (2010) Effect of solute stress and temperature on growth rate and *TRI5* gene expression using real time RT-PCR in *Fusarium graminearum* from Spanish wheat. *International Journal of Food Microbiology*, 140, 169–174.
- Matarese, F., Sarrocco, S., Gruber, S., Seidl-Seiboth, V. & Vannacci, G. (2012) Biocontrol of Fusarium head blight: interactions between *Trichoderma* and mycotoxigenic *Fusarium*. *Microbiology*, 158, 98–106.
- Matny, O.N. (2015) Fusarium head blight and crown rot on wheat & barley: losses and health risks. *Advances in Plants and Agriculture Research*, 2, 38–43.
- Merhej, J., Boutigny, A.L., Pinson-Gadais, L., Richard-Forget, F. & Barreau, C. (2010) Acidic pH as a determinant of *TRI* gene expression and trichothecene B biosynthesis in *Fusarium graminearum*. *Food Additives and Contaminants*, 27, 710–717.
- Merhej, J., Richard-Forget, F. & Barreau, C. (2011) The pH regulatory factor Pac1 regulates *tri* gene expression and trichothecene production in *Fusarium graminearum*. *Fungal Genetics and Biology*, 48, 275–284.
- Miedaner, T., Cumagun, C.J.R. & Chakraborty, S. (2008) Population genetics of three important head blight pathogens *Fusarium graminearum*, *F. pseudograminearum* and *F. culmorum*. *Journal of Phytopathology*, 156, 129–139.
- Mohamed, H.A.L.A. & Haggag, W.M. (2006) Biocontrol potential of salinity tolerant mutants of *Trichoderma harzianum* against *Fusarium oxysporum*. *Brazilian Journal of Microbiology*, 37, 181–191.
- Nagalakshmi, U., Waern, K. & Snyder, M. (2010) RNA-Seq: a method for comprehensive transcriptome analysis. *Current Protocols in Molecular Biology*, 89, 4–11.
- Nourozian, J., Etebarian, H.R. & Khodakaramian, G. (2006) Biological control of *Fusarium graminearum* on wheat by antagonistic bacteria. *Songklanakarinn Journal of Science and Technology*, 28(Suppl 1), 29–38.
- Oufensou, S., Balmes, V., Azara, E., Fabbri, D., Dettori, M.A., Schüller, C. et al. (2020) Naturally occurring phenols modulate vegetative growth and deoxynivalenol biosynthesis in *Fusarium graminearum*. *ACS Omega*, 5, 29407–29415.
- Özsoy, E., Kesercan, B. & Yörük, E. (2020) Antifungal activity of specific plant essential oils against *Fusarium graminearum*. *Phytopathology*, 90, 17–21.
- Pasquali, M. & Migheli, Q. (2014) Genetic approaches to chemotype determination in type B-trichothecene producing Fusaria. *International Journal of Food Microbiology*, 189, 164–182.
- Pertea, M., Pertea, G.M., Antonescu, C.M., Chang, T.C., Mendell, J.T. & Salzberg, S.L. (2015) StringTie enables improved reconstruction of a transcriptome from RNA-seq reads. *Nature Biotechnology*, 33, 290–295.
- Pinson-Gadais, L., Richard-Forget, F., Frasse, P., Barreau, C., Cahagnier, B., Richard-Molard, D. et al. (2008) Magnesium represses trichothecene biosynthesis and modulates *Tri5*, *Tri6*, and *Tri12* genes expression in *Fusarium graminearum*. *Mycopathologia*, 165, 51–59.
- Ponts, N., Pinson-Gadais, L., Barreau, C., Richard-Forget, F. & Ouellet, T. (2007) Exogenous H₂O₂ and catalase treatments interfere with *tri* genes expression in liquid cultures of *Fusarium graminearum*. *FEBS Letters*, 581, 443–447.
- Przemieniecki, S.W., Kurowski, T.P. & Korzekwa, K. (2014) Chemotypes and geographic distribution of the *Fusarium graminearum* species complex. *Environmental Biotechnology*, 10, 45–59.
- Ramangouda, G., Naik, M.K., Nitnavare, R.B., Yeshvekar, R., Bhattacharya, J., Bhatnagar-Mathur, P. et al. (2023) Genetic enhancement of *Trichoderma asperellum* biocontrol potentials and carbendazim tolerance for chickpea dry root rot disease management. *PLoS One*, 18, e0280064.
- Rawat, L., Singh, Y., Shukla, N. & Kumar, J. (2013) Salinity tolerant *Trichoderma harzianum* reinforces NaCl tolerance and reduces population dynamics of *Fusarium oxysporum* f. sp. *ciceri* in chickpea (*Cicer arietinum* L.) under salt stress conditions. *Archives of Phytopathology and Plant Protection*, 46, 1442–1467.

- Salmen, F., De Jonghe, J., Kaminski, T.S., Alemany, A., Parada, G.E., Verity-Legg, J. et al. (2022) High-throughput total RNA sequencing in single cells using VASA-seq. *Nature Biotechnology*, 40, 1780–1793.
- Sharma, V., Bhandari, P., Singh, B., Bhattacharya, A. & Shanmugam, V. (2013) Chitinase expression due to reduction in fusaric acid level in an antagonistic *Trichoderma harzianum* S17TH. *Indian Journal of Microbiology*, 53, 214–220.
- Sieber, C.M., Lee, W., Wong, P., Münsterkötter, M., Mewes, H.W., Schmeitzl, C. et al. (2014) The *Fusarium graminearum* genome reveals more secondary metabolite gene clusters and hints of horizontal gene transfer. *PLoS One*, 9, e110311.
- Strauch, O., Strasser, H., Hauschild, R. & Ehlers, R.U. (2011) Proposals for bacterial and fungal biocontrol agents. In: Ehlers, R.U. (Ed.) *Regulation of biological control agents*. Dordrecht: Springer, pp. 267–288.
- Waalwijk, C., Vanheule, A., Audenaert, K., Zhang, H., Warris, S., van de Geest, H. et al. (2017) *Fusarium* in the age of genomics. *Tropical Plant Pathology*, 42, 184–189.
- Wang, J.H., Ndoye, M., Zhang, J.B., Li, H.P. & Liao, Y.C. (2011) Population structure and genetic diversity of the *Fusarium graminearum* species complex. *Toxins*, 3, 1020–1037.
- Yaguchi, A., Yoshinari, T., Tsuyuki, R., Takahashi, H., Nakajima, T., Sugita-Konishi, Y. et al. (2009) Isolation and identification of precocenes and piperitone from essential oils as specific inhibitors of trichothecene production by *Fusarium graminearum*. *Journal of Agricultural and Food Chemistry*, 57, 846–851.
- Yang, Y., Li, M.X., Duan, Y.B., Li, T., Shi, Y.Y., Zhao, D.L. et al. (2018) A new point mutation in β 2-tubulin confers resistance to carbendazim in *Fusarium asiaticum*. *Pesticide Biochemistry and Physiology*, 145, 15–21.
- Yörük, E. (2018) Tetraconazole leads to alterations in *Fusarium graminearum* at different molecular levels. *Applied Ecology and Environment*, 16, 6155–6167.
- Yörük, E. & Albayrak, G. (2019) siRNA quelling of *tri4* and *tri5* genes related to deoxynivalenol synthesis in *Fusarium graminearum* and *Fusarium culmorum*. *Journal of Environmental Biology*, 40, 370–376.
- Yörük, E., Özkale, E., Sefer, Ö. & Özsoy, E. (2022) *Trichoderma atroviride* triggers alterations at epigenetics, transcripts, oxidative stress, and apoptosis levels on *Fusarium graminearum*. *Journal of Plant Pathology*, 104, 1039–1047.
- Yörük, E., Sefer, Ö., Tunali, B., Kansu, B., Sharifnabi, B. & Yli-Mattila, T. (2018) Thiophanate methyl susceptibility and alterations in *tri5*, *Mgv1* and *StuA* expression among *Fusarium graminearum* and *F. culmorum* isolates. *Journal of Plant Pathology*, 100, 447–455.
- Yörük, E. & Yli-Mattila, T. (2019) Class B-trichothecene profiles of *Fusarium* species as causal agents of head blight. In: Satyanarayana, T., Deshmukh, S. & Deshpande, M. (Eds.) *Advancing frontiers in mycology & mycotechnology*. Singapore: Springer, pp. 347–376.
- Zapparata, A., Baroncelli, R., Durling, M.B., Kubicek, C.P., Karlsson, M., Vannacci, G. et al. (2021) Fungal cross-talk: an integrated approach to study distance communication. *Fungal Genetics and Biology*, 148, 103518.
- Zeilinger, S., Gruber, S., Bansal, R. & Mukherjee, P.K. (2016) Secondary metabolism in *Trichoderma* – chemistry meets genomics. *Fungal Biology Reviews*, 30, 74–90.
- Zhao, C., Waalwijk, C., de Wit, P.J., Tang, D. & van der Lee, T. (2013) RNA-Seq analysis reveals new gene models and alternative splicing in the fungal pathogen *Fusarium graminearum*. *BMC Genomics*, 14, 21.
- Zheng, Z., Hou, Y., Cai, Y., Zhang, Y., Li, Y. & Zhou, M. (2015) Whole-genome sequencing reveals that mutations in *myosin-5* confer resistance to the fungicide phenamacril in *Fusarium graminearum*. *Scientific Reports*, 5, 8248.

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Özkale, E., Yörük, E., Budak, M. & Korkmaz, E.M. (2023) *Trichoderma atroviride* suppresses *Fusarium graminearum* by altering primary and secondary metabolite biosynthesis profiling. *Plant Pathology*, 72, 1428–1441. Available from: <https://doi.org/10.1111/ppa.13768>