

ASYMPTOTICAL INVARIANT AND ASYMPTOTICAL LACUNARY INVARIANT EQUIVALENCE TYPES FOR DOUBLE SEQUENCES VIA IDEALS USING MODULUS FUNCTIONS

ERDİNÇ DÜNDAR*, NİMET PANCAROĞLU AKIN, AND UĞUR ULUSU

Abstract. In this study, we present some asymptotical invariant and asymptotical lacunary invariant equivalence types for double sequences via ideals using modulus functions and investigate relationships between them.

1. Introduction and Background

The concept of convergence for double sequences was firstly introduced by Pringsheim [46]. Then, this concept was extended to the concept of statistical convergence by Mursaleen and Edely [31], to the concept of lacunary statistical convergence by Patterson and Savaş [43], to the concept of \mathcal{I}_2 -convergence by Das et al. [6] and to the concept of lacunary \mathcal{I}_2 -invariant convergence [56].

Patterson [42] introduced the concept of asymptotical equivalence for double sequences. Then, this concept was extended to the concept of asymptotical double statistical equivalence by Esi and Açıkgöz [13], to the concept of asymptotical double lacunary statistical equivalence by Esi [14], to the concept of asymptotical \mathcal{I}_2 -equivalence by Hazarika and Kumar [15] and to the concept of asymptotical lacunary \mathcal{I}_2 -invariant equivalence [57].

Modulus function was firstly introduced by Nakano [33]. Maddox [25], Pehlivan [45] and many other authors used a modulus function f to define new concepts and to give new theorems. Kumar and Sharma [23] studied on asymptotically lacunary equivalent sequences via ideals using modulus functions. For double sequences, the concept of f -asymptotical lacunary \mathcal{I}_2 -equivalence was studied by Dündar and P. Akın [10].

Received October 3, 2020. Revised February 11, 2021. Accepted February 15, 2021.
2020 Mathematics Subject Classification. 34C41, 40A05, 40A35.

Key words and phrases. Double sequence, Asymptotically equivalence, \mathcal{I} -convergence, Invariant convergence, Modulus function, Double lacunary sequence, Statistical convergence.

*Corresponding author

Several authors studied on the concepts of invariant mean, invariant convergence, asymptotical invariant equivalence of sequences and investigated some properties of these concepts (see [5, 9–12, 35, 39, 41, 47–49, 52, 56, 57]).

More study on the concepts of convergence or asymptotical equivalence for sequences can be found in [1–3, 7, 8, 16–21, 24, 26–32, 36–38, 40, 41, 44, 51, 54, 55, 58].

In this study, we present some asymptotical invariant and asymptotical lacunary invariant equivalence types for double sequences via ideals using modulus functions and investigate relationships between them.

Now, before giving the main part of the study, we recall some basic concepts (see [4, 9–11, 22, 25, 26, 34, 42, 43, 45, 50, 53]).

Two non-negative sequences $x = (x_{kj})$ and $y = (y_{kj})$ are said to be asymptotically equivalent if

$$\lim_{k,j \rightarrow \infty} \frac{x_{kj}}{y_{kj}} = 1$$

(denoted by $x \sim y$).

A family of sets $\mathcal{I} \subseteq 2^{\mathbb{N}}$ is called an ideal if and only if

(i) $\emptyset \in \mathcal{I}$, (ii) For each $A, B \in \mathcal{I}$ we have $A \cup B \in \mathcal{I}$, (iii) For each $A \in \mathcal{I}$ and each $B \subseteq A$ we have $B \in \mathcal{I}$.

An ideal $\mathcal{I} \subseteq 2^{\mathbb{N}}$ is called non-trivial if $\mathbb{N} \notin \mathcal{I}$ and a non-trivial ideal \mathcal{I} is called admissible if $\{n\} \in \mathcal{I}$, for each $n \in \mathbb{N}$.

A non-trivial ideal $\mathcal{I} \subseteq 2^{\mathbb{N} \times \mathbb{N}}$ is called strongly admissible ideal if $\{i\} \times \mathbb{N}$ and $\mathbb{N} \times \{i\}$ belong to \mathcal{I}_2 , for each $i \in \mathbb{N}$.

Throughout the study, we will consider the \mathcal{I}_2 as a strongly admissible ideal in $2^{\mathbb{N} \times \mathbb{N}}$.

It is evident that a strongly admissible ideal is admissible also.

Let $\mathcal{I}_2^0 = \{A \subset \mathbb{N} \times \mathbb{N} : (\exists m(A) \in \mathbb{N})(i, j \geq m(A) \Rightarrow (i, j) \notin A)\}$. Then, \mathcal{I}_2^0 is a strongly admissible ideal and clearly an ideal \mathcal{I}_2 is strongly admissible if and only if $\mathcal{I}_2^0 \subset \mathcal{I}_2$.

Two non-negative sequences $x = (x_{kj})$ and $y = (y_{kj})$ are said to be strong asymptotically \mathcal{I}_2 -equivalent of multiple L if for every $\varepsilon > 0$,

$$\left\{ (m, n) \in \mathbb{N} \times \mathbb{N} : \frac{1}{mn} \sum_{k,j=1,1}^{m,n} \left| \frac{x_{kj}}{y_{kj}} - L \right| \geq \varepsilon \right\} \in \mathcal{I}_2$$

(denoted by $x_{kj} \stackrel{[\mathcal{I}_2^L]}{\sim} y_{kj}$) and simply strong asymptotically \mathcal{I}_2 -equivalent if $L = 1$.

Two non-negative sequences $x = (x_{kj})$ and $y = (y_{kj})$ are said to be asymptotically \mathcal{I}_2 -statistical equivalent of multiple L if for every $\varepsilon > 0$ and each $\gamma > 0$,

$$\left\{ (m, n) \in \mathbb{N} \times \mathbb{N} : \frac{1}{mn} \left| \left\{ k, j \leq m, n : \left| \frac{x_{kj}}{y_{kj}} - L \right| \geq \varepsilon \right\} \right| \geq \gamma \right\} \in \mathcal{I}_2$$

(denoted by $x_{kj} \stackrel{\mathcal{I}_2^L(S)}{\sim} y_{kj}$) and simply asymptotically \mathcal{I}_2 -statistical equivalent if $L = 1$.

A function $f : [0, \infty) \rightarrow [0, \infty)$ is called a modulus if

1. $f(x) = 0$ if and only if $x = 0$,
2. $f(x + y) \leq f(x) + f(y)$,
3. f is increasing,
4. f is continuous from the right at 0.

A modulus may be unbounded (for example $f(x) = x^p$, $0 < p < 1$) or bounded (for example $f(x) = \frac{x}{x+1}$).

Let f be a modulus function. Two non-negative sequences $x = (x_{kj})$ and $y = (y_{kj})$ are said to be f -asymptotically \mathcal{I}_2 -equivalent of multiple L if for every $\varepsilon > 0$,

$$\left\{ (k, j) \in \mathbb{N} \times \mathbb{N} : f\left(\left|\frac{x_{kj}}{y_{kj}} - L\right|\right) \geq \varepsilon \right\} \in \mathcal{I}_2$$

(denoted by $x_{kj} \stackrel{\mathcal{I}_2^L(f)}{\sim} y_{kj}$) and simply f -asymptotically \mathcal{I}_2 -equivalent if $L = 1$.

Let f be a modulus function. Two non-negative sequences $x = (x_{kj})$ and $y = (y_{kj})$ are said to be strong f -asymptotically \mathcal{I}_2 -equivalent of multiple L if for every $\varepsilon > 0$,

$$\left\{ (m, n) \in \mathbb{N} \times \mathbb{N} : \frac{1}{mn} \sum_{k,j=1,1}^{m,n} f\left(\left|\frac{x_{kj}}{y_{kj}} - L\right|\right) \geq \varepsilon \right\} \in \mathcal{I}_2$$

(denoted by $x_{kj} \stackrel{[\mathcal{I}_2^L(f)]}{\sim} y_{kj}$) and simply strong f -asymptotically \mathcal{I}_2 -equivalent if $L = 1$.

The double sequence $\theta_2 = \{(k_r, j_u)\}$ is called double lacunary sequence if there exist two increasing sequence of integers such that

$$k_0 = 0, \quad h_r = k_r - k_{r-1} \rightarrow \infty \quad \text{and} \quad j_0 = 0, \quad \bar{h}_u = j_u - j_{u-1} \rightarrow \infty \quad \text{as } r, u \rightarrow \infty.$$

We use the following notations in the sequel:

$$h_{ru} = h_r \bar{h}_u, \quad I_{ru} = \{(k, j) : k_{r-1} < k \leq k_r \quad \text{and} \quad j_{u-1} < j \leq j_u\}.$$

Throughout the study, we will consider the $\theta_2 = \{(k_r, j_u)\}$ as a double lacunary sequence.

Two non-negative sequences $x = (x_{kj})$ and $y = (y_{kj})$ are said to be strong asymptotically lacunary \mathcal{I}_2 -equivalent of multiple L if for every $\varepsilon > 0$,

$$\left\{ (r, u) \in \mathbb{N} \times \mathbb{N} : \frac{1}{h_{ru}} \sum_{(k,j) \in I_{ru}} \left| \frac{x_{kj}}{y_{kj}} - L \right| \geq \varepsilon \right\} \in \mathcal{I}_2$$

(denoted by $x_{kj} \stackrel{[\mathcal{I}_{\theta_2}^L]}{\sim} y_{kj}$) and simply strong asymptotically lacunary \mathcal{I}_2 -equivalent if $L = 1$.

The two non-negative sequences $x = (x_{kj})$ and $y = (y_{kj})$ are said to be asymptotically lacunary \mathcal{I}_2 -statistical equivalent of multiple L if for every $\varepsilon > 0$

and each $\gamma > 0$,

$$\left\{ (r, u) \in \mathbb{N} \times \mathbb{N} : \frac{1}{h_{ru}} \left| \left\{ (k, j) \in I_{ru} : \left| \frac{x_{kj}}{y_{kj}} - L \right| \geq \varepsilon \right\} \right| \geq \gamma \right\} \in \mathcal{I}_2$$

(denoted by $x_{kj} \stackrel{\mathcal{I}_{\theta_2^L}(S)}{\sim} y_{kj}$) and simply asymptotically lacunary \mathcal{I}_2 -statistical equivalent if $L = 1$.

Let f be a modulus function. Two non-negative sequences $x = (x_{kj})$ and $y = (y_{kj})$ are said to be f -asymptotically lacunary \mathcal{I}_2 -equivalent of multiple L if for every $\varepsilon > 0$,

$$\left\{ (k, j) \in I_{ru} : f \left(\left| \frac{x_{kj}}{y_{kj}} - L \right| \right) \geq \varepsilon \right\} \in \mathcal{I}_2$$

(denoted by $x_{kj} \stackrel{\mathcal{I}_{\theta_2^L}(f)}{\sim} y_{kj}$) and simply f -asymptotically lacunary \mathcal{I}_2 -equivalent if $L = 1$.

Let f be a modulus function. Two non-negative sequences $x = (x_{kj})$ and $y = (y_{kj})$ are said to be strong f -asymptotically lacunary \mathcal{I}_2 -equivalent of multiple L if for every $\varepsilon > 0$,

$$\left\{ (r, u) \in \mathbb{N} \times \mathbb{N} : \frac{1}{h_{ru}} \sum_{(k,j) \in I_{ru}} f \left(\left| \frac{x_{kj}}{y_{kj}} - L \right| \right) \geq \varepsilon \right\} \in \mathcal{I}_2$$

(denoted by $x_{kj} \stackrel{[\mathcal{I}_{\theta_2^L}(f)]}{\sim} y_{kj}$) and simply strong f -asymptotically lacunary \mathcal{I}_2 -equivalent if $L = 1$.

Let σ be a mapping such that $\sigma : \mathbb{N} \rightarrow \mathbb{N}$. A continuous linear functional ψ on ℓ_∞ is said to be an invariant mean or a σ -mean if it satisfies the following conditions:

1. $\psi(x_n) \geq 0$, when the sequence (x_n) has $x_n \geq 0$ for all n ,
2. $\psi(e) = 1$, where $e = (1, 1, 1, \dots)$ and
3. $\psi(x_{\sigma(n)}) = \psi(x_n)$ for all $(x_n) \in \ell_\infty$.

The mappings σ are assumed to be one-to-one and such that $\sigma^m(n) \neq n$ for all $m, n \in \mathbb{N}$, where $\sigma^m(n)$ denotes the m th iterate of the mapping σ at n . Thus ψ extends the limit functional on c , in the sense that $\psi(x_n) = \lim x_n$ for all $(x_n) \in c$.

Let $E \subseteq \mathbb{N} \times \mathbb{N}$ and

$$s_{mk} = \min_{i,j} \left| E \cap \left\{ (\sigma(i), \sigma(j)), (\sigma^2(i), \sigma^2(j)), \dots, (\sigma^m(i), \sigma^k(j)) \right\} \right|$$

and

$$S_{mk} = \max_{i,j} \left| E \cap \left\{ (\sigma(i), \sigma(j)), (\sigma^2(i), \sigma^2(j)), \dots, (\sigma^m(i), \sigma^k(j)) \right\} \right|.$$

If the limits $\underline{V}_2(E) = \lim_{m,k \rightarrow \infty} \frac{s_{mk}}{mk}$ and $\overline{V}_2(E) = \lim_{m,k \rightarrow \infty} \frac{S_{mk}}{mk}$ exists, then they are called a lower σ -uniform density and an upper σ -uniform density of

the set E , respectively. If $\underline{V}_2(E) = \overline{V}_2(E)$, then $V_2(E) = \underline{V}_2(E) = \overline{V}_2(E)$ is called σ -uniform density of the set E .

Denote by \mathcal{I}_2^σ the class of all $E \subseteq \mathbb{N} \times \mathbb{N}$ with $V_2(E) = 0$. Obviously \mathcal{I}_2^σ is a strongly admissible ideal in $\mathbb{N} \times \mathbb{N}$.

Two non-negative sequences $x = (x_{kj})$ and $y = (y_{kj})$ are said to be asymptotically \mathcal{I}_2 -invariant equivalent of multiple L if for every $\varepsilon > 0$,

$$\left\{ (k, j) \in \mathbb{N} \times \mathbb{N} : \left| \frac{x_{kj}}{y_{kj}} - L \right| \geq \varepsilon \right\} \in \mathcal{I}_2^\sigma$$

(denoted by $x_{kj} \stackrel{\mathcal{I}_2^\sigma}{\sim} y_{kj}$) and simply asymptotically \mathcal{I}_2 -invariant equivalent if $L = 1$.

Let $\theta_2 = \{(k_r, j_u)\}$ be a double lacunary sequence, $C \subseteq \mathbb{N} \times \mathbb{N}$ and

$$s_{ru} = \min_{m,n} \left| C \cap \left\{ (\sigma^k(m), \sigma^j(n)) : (k, j) \in I_{ru} \right\} \right|$$

and

$$S_{ru} = \max_{m,n} \left| C \cap \left\{ (\sigma^k(m), \sigma^j(n)) : (k, j) \in I_{ru} \right\} \right|.$$

If the limits $\underline{V}_2^\theta(C) = \lim_{r,u \rightarrow \infty} \frac{s_{ru}}{h_{ru}}$ and $\overline{V}_2^\theta(C) = \lim_{r,u \rightarrow \infty} \frac{S_{ru}}{h_{ru}}$ exist, then they are called a lower lacunary σ -uniform density and an upper lacunary σ -uniform density of the set C , respectively. If $\underline{V}_2^\theta(C) = \overline{V}_2^\theta(C)$, then $V_2^\theta(C) = \underline{V}_2^\theta(C) = \overline{V}_2^\theta(C)$ is called the lacunary σ -uniform density of the set C .

Denote by $\mathcal{I}_2^{\sigma\theta}$ the class of all $C \subseteq \mathbb{N} \times \mathbb{N}$ with $V_2^\theta(C) = 0$. Obviously $\mathcal{I}_2^{\sigma\theta}$ is a strongly admissible ideal in $\mathbb{N} \times \mathbb{N}$.

Two non-negative sequences $x = (x_{kj})$ and $y = (y_{kj})$ are said to be asymptotically lacunary \mathcal{I}_2 -invariant equivalent of multiple L if for every $\varepsilon > 0$,

$$\left\{ (k, j) \in I_{ru} : \left| \frac{x_{kj}}{y_{kj}} - L \right| \geq \varepsilon \right\} \in \mathcal{I}_2^{\sigma\theta}$$

(denoted by $x_{kj} \stackrel{\mathcal{I}_2^{\sigma\theta}}{\sim} y_{kj}$) and simply asymptotically lacunary \mathcal{I}_2 -invariant equivalent if $L = 1$.

Lemma 1.1. [45] *Let f be a modulus and $0 < \delta < 1$. Then, for each $x \geq \delta$ we have $f(x) \leq 2f(1)\delta^{-1}x$.*

2. MAIN RESULTS

Definition 2.1. *Two non-negative sequences $x = (x_{kj})$ and $y = (y_{kj})$ are strong asymptotically \mathcal{I}_2 -invariant equivalent of multiple L if for every $\varepsilon > 0$,*

$$\left\{ (m, n) \in \mathbb{N} \times \mathbb{N} : \frac{1}{mn} \sum_{k,j=1,1}^{m,n} \left| \frac{x_{kj}}{y_{kj}} - L \right| \geq \varepsilon \right\} \in \mathcal{I}_2^\sigma$$

(denoted by $x_{kj} \stackrel{[\mathcal{I}_{\sigma_2}^L]}{\sim} y_{kj}$) and simply strong asymptotically \mathcal{I}_2 -invariant equivalent if $L = 1$.

Definition 2.2. Let f be a modulus function. Two non-negative sequences $x = (x_{kj})$ and $y = (y_{kj})$ are f -asymptotically \mathcal{I}_2 -invariant equivalent of multiple L if for every $\varepsilon > 0$,

$$\left\{ (k, j) \in \mathbb{N} \times \mathbb{N} : f\left(\left|\frac{x_{kj}}{y_{kj}} - L\right|\right) \geq \varepsilon \right\} \in \mathcal{I}_2^\sigma$$

(denoted by $x_{kj} \stackrel{\mathcal{I}_{\sigma_2}^L(f)}{\sim} y_{kj}$) and simply f -asymptotically \mathcal{I}_2 -invariant equivalent if $L = 1$.

Definition 2.3. Let f be a modulus function. Two non-negative sequences $x = (x_{kj})$ and $y = (y_{kj})$ are strong f -asymptotically \mathcal{I}_2 -invariant equivalent of multiple L if for every $\varepsilon > 0$,

$$\left\{ (m, n) \in \mathbb{N} \times \mathbb{N} : \frac{1}{mn} \sum_{k,j=1,1}^{m,n} f\left(\left|\frac{x_{kj}}{y_{kj}} - L\right|\right) \geq \varepsilon \right\} \in \mathcal{I}_2^\sigma$$

(denoted by $x_{kj} \stackrel{[\mathcal{I}_{\sigma_2}^L(f)]}{\sim} y_{kj}$) and simply strong f -asymptotically \mathcal{I}_2 -invariant equivalent if $L = 1$.

Theorem 2.4. Let f be a modulus function. Then,

$$x_{kj} \stackrel{[\mathcal{I}_{\sigma_2}^L]}{\sim} y_{kj} \Rightarrow x_{kj} \stackrel{[\mathcal{I}_{\sigma_2}^L(f)]}{\sim} y_{kj}.$$

Proof. Suppose that $x_{kj} \stackrel{[\mathcal{I}_{\sigma_2}^L]}{\sim} y_{kj}$ and $\varepsilon > 0$ be given. Choose $0 < \delta < 1$ such that $f(t) < \varepsilon$, for $0 \leq t \leq \delta$. We can write

$$\begin{aligned} \frac{1}{mn} \sum_{k,j=1,1}^{m,n} f\left(\left|\frac{x_{kj}}{y_{kj}} - L\right|\right) &= \frac{1}{mn} \sum_{k,j=1,1}^{m,n} f\left(\left|\frac{x_{kj}}{y_{kj}} - L\right|\right) \\ &\quad \left|\frac{x_{kj}}{y_{kj}} - L\right| \leq \delta \\ &+ \frac{1}{mn} \sum_{k,j=1,1}^{m,n} f\left(\left|\frac{x_{kj}}{y_{kj}} - L\right|\right) \\ &\quad \left|\frac{x_{kj}}{y_{kj}} - L\right| > \delta \end{aligned}$$

and so, by Lemma 1.1, we have

$$\frac{1}{mn} \sum_{k,j=1,1}^{m,n} f\left(\left|\frac{x_{kj}}{y_{kj}} - L\right|\right) < \varepsilon + \left(\frac{2f(1)}{\delta}\right) \frac{1}{mn} \sum_{k,j=1,1}^{m,n} \left|\frac{x_{kj}}{y_{kj}} - L\right|.$$

Thus, for any $\gamma > 0$

$$\begin{aligned} & \left\{ (m, n) \in \mathbb{N} \times \mathbb{N} : \frac{1}{mn} \sum_{k,j=1,1}^{m,n} f\left(\left|\frac{x_{kj}}{y_{kj}} - L\right|\right) \geq \gamma \right\} \\ & \subseteq \left\{ (m, n) \in \mathbb{N} \times \mathbb{N} : \frac{1}{mn} \sum_{k,j=1,1}^{m,n} \left|\frac{x_{kj}}{y_{kj}} - L\right| \geq \frac{(\gamma - \varepsilon)\delta}{2f(1)} \right\}. \end{aligned}$$

Since it is $x_{kj} \stackrel{[\mathcal{I}_{\sigma_2}^L]}{\sim} y_{kj}$, the second set in the above expression and therefore the first set also belongs to the \mathcal{I}_2^σ . This proves that $x_{kj} \stackrel{[\mathcal{I}_{\sigma_2}^L(f)]}{\sim} y_{kj}$. \square

Theorem 2.5. *If $\lim_{t \rightarrow \infty} \frac{f(t)}{t} = \alpha > 0$, then*

$$x_{kj} \stackrel{[\mathcal{I}_{\sigma_2}^L(f)]}{\sim} y_{kj} \Leftrightarrow x_{kj} \stackrel{[\mathcal{I}_{\sigma_2}^L]}{\sim} y_{kj}.$$

Proof. We showed that $x_{kj} \stackrel{[\mathcal{I}_{\sigma_2}^L]}{\sim} y_{kj} \Rightarrow x_{kj} \stackrel{[\mathcal{I}_{\sigma_2}^L(f)]}{\sim} y_{kj}$ in Theorem 2.4. Now, for proof, we must show $x_{kj} \stackrel{[\mathcal{I}_{\sigma_2}^L(f)]}{\sim} y_{kj} \Rightarrow x_{kj} \stackrel{[\mathcal{I}_{\sigma_2}^L]}{\sim} y_{kj}$.

Assume that $x_{kj} \stackrel{[\mathcal{I}_{\sigma_2}^L(f)]}{\sim} y_{kj}$. Let $\lim_{t \rightarrow \infty} \frac{f(t)}{t} = \alpha > 0$, then $f(t) \geq \alpha t$ for all $t \geq 0$. So, we can write

$$\begin{aligned} \frac{1}{mn} \sum_{k,j=1,1}^{m,n} f\left(\left|\frac{x_{kj}}{y_{kj}} - L\right|\right) & \geq \frac{1}{mn} \sum_{k,j=1,1}^{m,n} \alpha \left(\left|\frac{x_{kj}}{y_{kj}} - L\right|\right) \\ & = \alpha \left(\frac{1}{mn} \sum_{k,j=1,1}^{m,n} \left|\frac{x_{kj}}{y_{kj}} - L\right|\right), \end{aligned}$$

and it follows that for every $\varepsilon > 0$, we have

$$\begin{aligned} & \left\{ (m, n) \in \mathbb{N} \times \mathbb{N} : \frac{1}{mn} \sum_{k,j=1,1}^{m,n} \left|\frac{x_{kj}}{y_{kj}} - L\right| \geq \varepsilon \right\} \\ & \subseteq \left\{ (m, n) \in \mathbb{N} \times \mathbb{N} : \frac{1}{mn} \sum_{k,j=1,1}^{m,n} f\left(\left|\frac{x_{kj}}{y_{kj}} - L\right|\right) \geq \alpha\varepsilon \right\}. \end{aligned}$$

Since it is $x_{kj} \stackrel{[\mathcal{I}_{\sigma_2}^L(f)]}{\sim} y_{kj}$, the second set in the above expression and therefore the first set also belongs to the \mathcal{I}_2^σ . This proves that $x_{kj} \stackrel{[\mathcal{I}_{\sigma_2}^L(f)]}{\sim} y_{kj} \Rightarrow x_{kj} \stackrel{[\mathcal{I}_{\sigma_2}^L]}{\sim} y_{kj}$ and so $x_{kj} \stackrel{[\mathcal{I}_{\sigma_2}^L(f)]}{\sim} y_{kj} \Leftrightarrow x_{kj} \stackrel{[\mathcal{I}_{\sigma_2}^L]}{\sim} y_{kj}$. \square

Definition 2.6. *Two non-negative sequences $x = (x_{kj})$ and $y = (y_{kj})$ are asymptotically \mathcal{I}_2 -invariant statistical equivalent of multiple L if for every $\varepsilon > 0$*

and each $\gamma > 0$,

$$\left\{ (m, n) \in \mathbb{N} \times \mathbb{N} : \frac{1}{mn} \left| \left\{ k, j \leq m, n : \left| \frac{x_{kj}}{y_{kj}} - L \right| \geq \varepsilon \right\} \right| \geq \gamma \right\} \in \mathcal{I}_2^\sigma$$

(denoted by $x_{kj} \stackrel{\mathcal{I}_{\sigma_2}^L(S)}{\sim} y_{kj}$) and simply asymptotically \mathcal{I}_2 -invariant statistical equivalent if $L = 1$.

Theorem 2.7. *Let f be a modulus function. Then,*

$$x_{kj} \stackrel{[\mathcal{I}_{\sigma_2}^L(f)]}{\sim} y_{kj} \Rightarrow x_{kj} \stackrel{\mathcal{I}_{\sigma_2}^L(S)}{\sim} y_{kj}.$$

Proof. Assume that $x_{kj} \stackrel{[\mathcal{I}_{\sigma_2}^L(f)]}{\sim} y_{kj}$ and $\varepsilon > 0$ be given. We can write

$$\begin{aligned} \frac{1}{mn} \sum_{k,j=1,1}^{m,n} f\left(\left|\frac{x_{kj}}{y_{kj}} - L\right|\right) &\geq \frac{1}{mn} \sum_{k,j=1,1}^{m,n} f\left(\left|\frac{x_{kj}}{y_{kj}} - L\right|\right) \\ &\quad \left|\frac{x_{kj}}{y_{kj}} - L\right| \geq \varepsilon \\ &\geq f(\varepsilon) \frac{1}{mn} \left| \left\{ k \leq m, j \leq n : \left| \frac{x_{kj}}{y_{kj}} - L \right| \geq \varepsilon \right\} \right|. \end{aligned}$$

Thus, for any $\gamma > 0$ we have

$$\begin{aligned} &\left\{ (m, n) \in \mathbb{N} \times \mathbb{N} : \frac{1}{mn} \left| \left\{ k \leq m, j \leq n : \left| \frac{x_{kj}}{y_{kj}} - L \right| \geq \varepsilon \right\} \right| \geq \frac{\gamma}{f(\varepsilon)} \right\} \\ &\subseteq \left\{ (m, n) \in \mathbb{N} \times \mathbb{N} : \frac{1}{mn} \sum_{k,j=1,1}^{m,n} f\left(\left|\frac{x_{kj}}{y_{kj}} - L\right|\right) \geq \gamma \right\}. \end{aligned}$$

Since it is $x_{kj} \stackrel{[\mathcal{I}_{\sigma_2}^L(f)]}{\sim} y_{kj}$, the second set in the above expression and therefore the first set also belongs to the \mathcal{I}_2^σ . This proves that $x_{kj} \stackrel{\mathcal{I}_{\sigma_2}^L(S)}{\sim} y_{kj}$. \square

Theorem 2.8. *Let f be a modulus function. If f is bounded, then*

$$x_{kj} \stackrel{\mathcal{I}_{\sigma_2}^L(S)}{\sim} y_{kj} \Leftrightarrow x_{kj} \stackrel{[\mathcal{I}_{\sigma_2}^L(f)]}{\sim} y_{kj}.$$

Proof. We showed that $x_{kj} \stackrel{[\mathcal{I}_{\sigma_2}^L(f)]}{\sim} y_{kj} \Rightarrow x_{kj} \stackrel{\mathcal{I}_{\sigma_2}^L(S)}{\sim} y_{kj}$ in Theorem 2.7.

Now, for proof, we must show $x_{kj} \stackrel{\mathcal{I}_{\sigma_2}^L(S)}{\sim} y_{kj} \Rightarrow x_{kj} \stackrel{[\mathcal{I}_{\sigma_2}^L(f)]}{\sim} y_{kj}$.

Assume that $x_{kj} \stackrel{\mathcal{I}_{\sigma_2}^L(S)}{\sim} y_{kj}$. Let f is bounded, then there exists a positive real number M such that $|f(x)| \leq M$, for all $x \geq 0$. So, we have

$$\begin{aligned} \frac{1}{mn} \sum_{k,j=1,1}^{m,n} f\left(\left|\frac{x_{kj}}{y_{kj}} - L\right|\right) &= \frac{1}{mn} \sum_{k,j=1,1}^{m,n} f\left(\left|\frac{x_{kj}}{y_{kj}} - L\right|\right) \\ &\quad \left|\frac{x_{kj}}{y_{kj}} - L\right| \geq \varepsilon \\ &+ \frac{1}{mn} \sum_{k,j=1,1}^{m,n} f\left(\left|\frac{x_{kj}}{y_{kj}} - L\right|\right) \\ &\quad \left|\frac{x_{kj}}{y_{kj}} - L\right| < \varepsilon \\ &\leq \frac{M}{mn} \left| \left\{ k \leq m, j \leq n : \left|\frac{x_{kj}}{y_{kj}} - L\right| \geq \varepsilon \right\} \right| \\ &\quad + f(\varepsilon). \end{aligned}$$

This proves that $x_{kj} \stackrel{[\mathcal{I}_{\sigma_2}^L(f)]}{\sim} y_{kj}$. \square

Definition 2.9. Two non-negative sequences $x = (x_{kj})$ and $y = (y_{kj})$ are strong asymptotically lacunary \mathcal{I}_2 -invariant equivalent of multiple L if for every $\varepsilon > 0$,

$$\left\{ (r, u) \in \mathbb{N} \times \mathbb{N} : \frac{1}{h_{ru}} \sum_{(k,j) \in I_{ru}} \left| \frac{x_{kj}}{y_{kj}} - L \right| \geq \varepsilon \right\} \in \mathcal{I}_2^\theta$$

(denoted by $x_{kj} \stackrel{[\mathcal{I}_{\sigma_2}^{\theta L}]}{\sim} y_{kj}$) and simply strong asymptotically lacunary \mathcal{I}_2 -invariant equivalent if $L = 1$.

Definition 2.10. Let f be a modulus function. Two non-negative sequences $x = (x_{kj})$ and $y = (y_{kj})$ are f -asymptotically lacunary \mathcal{I}_2 -invariant equivalent of multiple L if for every $\varepsilon > 0$,

$$\left\{ (k, j) \in I_{ru} : f\left(\left|\frac{x_{kj}}{y_{kj}} - L\right|\right) \geq \varepsilon \right\} \in \mathcal{I}_2^\theta$$

(denoted by $x_{kj} \stackrel{\mathcal{I}_{\sigma_2}^{\theta L}(f)}{\sim} y_{kj}$) and simply f -asymptotically lacunary \mathcal{I}_2 -invariant equivalent if $L = 1$.

Definition 2.11. Let f be a modulus function. Two non-negative sequences $x = (x_{kj})$ and $y = (y_{kj})$ are strong f -asymptotically lacunary \mathcal{I}_2 -invariant equivalent of multiple L if for every $\varepsilon > 0$,

$$\left\{ (r, u) \in \mathbb{N} \times \mathbb{N} : \frac{1}{h_{ru}} \sum_{(k,j) \in I_{ru}} f\left(\left|\frac{x_{kj}}{y_{kj}} - L\right|\right) \geq \varepsilon \right\} \in \mathcal{I}_2^{\sigma\theta}$$

(denoted by $x_{kj} \stackrel{[\mathcal{I}_{\sigma_2}^{\theta L}(f)]}{\sim} y_{kj}$) and simply strong f -asymptotically lacunary \mathcal{I}_2 -invariant equivalent if $L = 1$.

Theorem 2.12. *Let f be a modulus function. Then,*

$$x_{kj} \stackrel{[\mathcal{I}_{\sigma_2}^{\theta L}]}{\sim} y_{kj} \Rightarrow x_{kj} \stackrel{[\mathcal{I}_{\sigma_2}^{\theta L}(f)]}{\sim} y_{kj}.$$

Proof. Suppose that $x_{kj} \stackrel{[\mathcal{I}_{\sigma_2}^{\theta L}]}{\sim} y_{kj}$ and $\varepsilon > 0$ be given. Choose $0 < \delta < 1$ such that $f(t) < \varepsilon$, for $0 \leq t \leq \delta$. We can write

$$\begin{aligned} \frac{1}{h_{ru}} \sum_{(k,j) \in I_{ru}} f\left(\left|\frac{x_{kj}}{y_{kj}} - L\right|\right) &= \frac{1}{h_{ru}} \sum_{(k,j) \in I_{ru}} f\left(\left|\frac{x_{kj}}{y_{kj}} - L\right|\right) \\ &\quad \left|\frac{x_{kj}}{y_{kj}} - L\right| \leq \delta \\ &\quad + \frac{1}{h_{ru}} \sum_{(k,j) \in I_{ru}} f\left(\left|\frac{x_{kj}}{y_{kj}} - L\right|\right) \\ &\quad \left|\frac{x_{kj}}{y_{kj}} - L\right| > \delta \end{aligned}$$

and so, by Lemma 1.1, we have

$$\frac{1}{h_{ru}} \sum_{(k,j) \in I_{ru}} f\left(\left|\frac{x_{kj}}{y_{kj}} - L\right|\right) < \varepsilon + \left(\frac{2f(1)}{\delta}\right) \frac{1}{h_{ru}} \sum_{(k,j) \in I_{ru}} \left|\frac{x_{kj}}{y_{kj}} - L\right|.$$

Thus, for any $\gamma > 0$

$$\begin{aligned} &\left\{ (r, u) \in \mathbb{N} \times \mathbb{N} : \frac{1}{h_{ru}} \sum_{(k,j) \in I_{ru}} f\left(\left|\frac{x_{kj}}{y_{kj}} - L\right|\right) \geq \gamma \right\} \\ &\subseteq \left\{ (r, u) \in \mathbb{N} \times \mathbb{N} : \frac{1}{h_{ru}} \sum_{(k,j) \in I_{ru}} \left|\frac{x_{kj}}{y_{kj}} - L\right| \geq \frac{(\gamma - \varepsilon)\delta}{2f(1)} \right\}. \end{aligned}$$

Since it is $x_{kj} \stackrel{[\mathcal{I}_{\sigma_2}^{\theta L}]}{\sim} y_{kj}$, the second set in the above expression and therefore the first set also belongs to the $\mathcal{I}_2^{\theta\theta}$. This proves that $x_{kj} \stackrel{[\mathcal{I}_{\sigma_2}^{\theta L}(f)]}{\sim} y_{kj}$. \square

Theorem 2.13. *If $\lim_{t \rightarrow \infty} \frac{f(t)}{t} = \alpha > 0$, then*

$$x_{kj} \stackrel{[\mathcal{I}_{\sigma_2}^{\theta L}(f)]}{\sim} y_{kj} \Leftrightarrow x_{kj} \stackrel{[\mathcal{I}_{\sigma_2}^{\theta L}]}{\sim} y_{kj}.$$

Proof. We showed that $x_{kj} \stackrel{[\mathcal{I}_{\sigma_2}^{\theta L}]}{\sim} y_{kj} \Rightarrow x_{kj} \stackrel{[\mathcal{I}_{\sigma_2}^{\theta L}(f)]}{\sim} y_{kj}$ in Theorem 2.12. Now, for proof, we must show $x_{kj} \stackrel{[\mathcal{I}_{\sigma_2}^{\theta L}(f)]}{\sim} y_{kj} \Rightarrow x_{kj} \stackrel{[\mathcal{I}_{\sigma_2}^{\theta L}]}{\sim} y_{kj}$.

Assume that $x_{kj} \stackrel{[\mathcal{I}_{\sigma_2}^{\theta L}(f)]}{\sim} y_{kj}$. Let $\lim_{t \rightarrow \infty} \frac{f(t)}{t} = \alpha > 0$, then $f(t) \geq \alpha t$ for all $t \geq 0$. So, we can write

$$\begin{aligned} \frac{1}{h_{ru}} \sum_{(k,j) \in I_{ru}} f\left(\left|\frac{x_{kj}}{y_{kj}} - L\right|\right) &\geq \frac{1}{h_{ru}} \sum_{(k,j) \in I_{ru}} \alpha \left(\left|\frac{x_{kj}}{y_{kj}} - L\right|\right) \\ &= \alpha \left(\frac{1}{h_{ru}} \sum_{(k,j) \in I_{ru}} \left|\frac{x_{kj}}{y_{kj}} - L\right|\right), \end{aligned}$$

and it follows that for every $\varepsilon > 0$, we have

$$\begin{aligned} &\left\{ (r, u) \in \mathbb{N} \times \mathbb{N} : \frac{1}{h_{ru}} \sum_{(k,j) \in I_{ru}} \left|\frac{x_{kj}}{y_{kj}} - L\right| \geq \varepsilon \right\} \\ &\subseteq \left\{ (r, u) \in \mathbb{N} \times \mathbb{N} : \frac{1}{h_{ru}} \sum_{(k,j) \in I_{ru}} f\left(\left|\frac{x_{kj}}{y_{kj}} - L\right|\right) \geq \alpha \varepsilon \right\}. \end{aligned}$$

Since it is $x_{kj} \stackrel{[\mathcal{I}_{\sigma_2}^{\theta L}(f)]}{\sim} y_{kj}$, the second set in the above expression and therefore the first set also belongs to the $\mathcal{I}_2^{\sigma\theta}$. This proves that $x_{kj} \stackrel{[\mathcal{I}_{\sigma_2}^{\theta L}(f)]}{\sim} y_{kj} \Rightarrow x_{kj} \stackrel{[\mathcal{I}_{\sigma_2}^{\theta L}]}{\sim} y_{kj}$ and so $x_{kj} \stackrel{[\mathcal{I}_{\sigma_2}^{\theta L}(f)]}{\sim} y_{kj} \Leftrightarrow x_{kj} \stackrel{[\mathcal{I}_{\sigma_2}^{\theta L}]}{\sim} y_{kj}$. \square

Definition 2.14. Two non-negative sequences $x = (x_{kj})$ and $y = (y_{kj})$ are asymptotically lacunary \mathcal{I}_2 -invariant statistical equivalent of multiple L if for every $\varepsilon > 0$ and each $\gamma > 0$,

$$\left\{ (r, u) \in \mathbb{N} \times \mathbb{N} : \frac{1}{h_{ru}} \left| \left\{ (k, j) \in I_{ru} : \left|\frac{x_{kj}}{y_{kj}} - L\right| \geq \varepsilon \right\} \right| \geq \gamma \right\} \in \mathcal{I}_2^{\sigma\theta}$$

(denoted by $x_{kj} \stackrel{\mathcal{I}_{\sigma_2}^{\theta L}(S)}{\sim} y_{kj}$) and simply asymptotically lacunary \mathcal{I}_2 -invariant statistical equivalent if $L = 1$.

Theorem 2.15. Let f be a modulus function. Then,

$$x_{kj} \stackrel{[\mathcal{I}_{\sigma_2}^{\theta L}(f)]}{\sim} y_{kj} \Rightarrow x_{kj} \stackrel{\mathcal{I}_{\sigma_2}^{\theta L}(S)}{\sim} y_{kj}.$$

Proof. Assume that $x_{kj} \stackrel{[\mathcal{I}_{\sigma_2}^{\theta L}(f)]}{\sim} y_{kj}$ and $\varepsilon > 0$ be given. We can write

$$\begin{aligned} \frac{1}{h_{ru}} \sum_{(k,j) \in I_{ru}} f\left(\left|\frac{x_{kj}}{y_{kj}} - L\right|\right) &\geq \frac{1}{h_{ru}} \sum_{\substack{(k,j) \in I_{ru} \\ \left|\frac{x_{kj}}{y_{kj}} - L\right| \geq \varepsilon}} f\left(\left|\frac{x_{kj}}{y_{kj}} - L\right|\right) \\ &\geq f(\varepsilon) \frac{1}{h_{ru}} \left| \left\{ (k, j) \in I_{ru} : \left|\frac{x_{kj}}{y_{kj}} - L\right| \geq \varepsilon \right\} \right|. \end{aligned}$$

Thus, for any $\gamma > 0$, we have

$$\begin{aligned} & \left\{ (r, u) \in \mathbb{N} \times \mathbb{N} : \frac{1}{h_{ru}} \left| \left\{ (k, j) \in I_{ru} : \left| \frac{x_{kj}}{y_{kj}} - L \right| \geq \varepsilon \right\} \right| \geq \gamma \right\} \\ & \subseteq \left\{ (r, u) \in \mathbb{N} \times \mathbb{N} : \frac{1}{h_{ru}} \sum_{(k,j) \in I_{ru}} f \left(\left| \frac{x_{kj}}{y_{kj}} - L \right| \right) \geq \gamma f(\varepsilon) \right\}. \end{aligned}$$

Since it is $x_{kj} \overset{[\mathcal{I}_{\sigma_2}^{\theta L}(f)]}{\sim} y_{kj}$, the second set in the above expression and therefore the first set also belongs to the $\mathcal{I}_2^{\sigma\theta}$. This proves that $x_{kj} \overset{[\mathcal{I}_{\sigma_2}^{\theta L}(S)]}{\sim} y_{kj}$. \square

Theorem 2.16. *Let f be a modulus function. If f is bounded, then*

$$x_{kj} \overset{[\mathcal{I}_{\sigma_2}^{\theta L}(S)]}{\sim} y_{kj} \Leftrightarrow x_{kj} \overset{[\mathcal{I}_{\sigma_2}^{\theta L}(f)]}{\sim} y_{kj}.$$

Proof. We showed that $x_{kj} \overset{[\mathcal{I}_{\sigma_2}^{\theta L}(f)]}{\sim} y_{kj} \Rightarrow x_{kj} \overset{[\mathcal{I}_{\sigma_2}^{\theta L}(S)]}{\sim} y_{kj}$ in Theorem 2.15.

Now, for proof, we must show $x_{kj} \overset{[\mathcal{I}_{\sigma_2}^{\theta L}(S)]}{\sim} y_{kj} \Rightarrow x_{kj} \overset{[\mathcal{I}_{\sigma_2}^{\theta L}(f)]}{\sim} y_{kj}$.

Assume that $x_{kj} \overset{[\mathcal{I}_{\sigma_2}^{\theta L}(S)]}{\sim} y_{kj}$. Let f is bounded, then there exists a positive real number M such that $|f(x)| \leq M$, for all $x \geq 0$. So, we have

$$\begin{aligned} \frac{1}{h_{ru}} \sum_{(k,j) \in I_{ru}} f \left(\left| \frac{x_{kj}}{y_{kj}} - L \right| \right) &= \frac{1}{h_{ru}} \sum_{(k,j) \in I_{ru}} f \left(\left| \frac{x_{kj}}{y_{kj}} - L \right| \right) \\ & \quad \left| \frac{x_{kj}}{y_{kj}} - L \right| \geq \varepsilon \\ &+ \frac{1}{h_{ru}} \sum_{(k,j) \in I_{ru}} f \left(\left| \frac{x_{kj}}{y_{kj}} - L \right| \right) \\ & \quad \left| \frac{x_{kj}}{y_{kj}} - L \right| < \varepsilon \\ &\leq \frac{M}{h_{ru}} \left| \left\{ (k, j) \in I_{ru} : \left| \frac{x_{kj}}{y_{kj}} - L \right| \geq \varepsilon \right\} \right| \\ &+ f(\varepsilon). \end{aligned}$$

This proves that $x_{kj} \overset{[\mathcal{I}_{\sigma_2}^{\theta L}(f)]}{\sim} y_{kj}$. \square

References

- [1] M. Arslan, E. Dündar, *On \mathcal{I} -convergence of sequences of functions in 2-normed spaces*, Southeast Asian Bull. Math. **42** (2018), 491–502.
- [2] N.L. Braha, H.M. Srivastava, S.A. Mohiuddine, *A Korovkin's type approximation theorem for periodic functions via the statistical summability of the generalized de la Vallée Poussin mean*, Appl. Math. Comput. **228** (2014), 162–169.

- [3] C. Belen, S.A. Mohiuddine, *Generalized weighted statistical convergence and application*, Appl. Math. Comput. **219** (2013), 9821–9826
- [4] J.S. Connor, *On strong matrix summability with respect to a modulus and statistical convergence*, Canad. Math. Bull. **32** (1989), 194–198.
- [5] C. Çakan, B. Altay, and M. Mursaleen, *The σ -convergence and σ -core of double sequences*, Appl. Math. Lett. **19** (2006), 1122–1128.
- [6] P. Das, P. Kostyrko, W. Wilczyński, P. Malik, *\mathcal{I} and \mathcal{I}^* -convergence of double sequences*, Math. Slovaca, **58**(5) (2008), 605–620.
- [7] E. Dündar, B. Altay, *\mathcal{I}_2 -convergence and \mathcal{I}_2 -Cauchy of double sequences*, Acta Math. Sci. **34B**(2) (2014), 343–353.
- [8] E. Dündar, B. Altay, *\mathcal{I}_2 -uniform convergence of double sequences of functions*, Filomat, **30**(5) (2016), 1273–1281.
- [9] E. Dündar, N. Pancaroğlu Akın, *f -asymptotically \mathcal{I}_σ -equivalence of real sequences*, Konuralp J. Math. **8**(1) (2020), 207–210
- [10] E. Dündar, N. Pancaroğlu Akın, *f -asymptotically $\mathcal{I}_{\sigma\theta}$ -equivalence of real sequences*, J. Math. Sci. Model. **3**(1) (2020), 32–37.
- [11] E. Dündar, N. Pancaroğlu Akın, *Wijsman lacunary ideal invariant convergence of double sequences of sets*, Honam Math. J. **42**(2) (2020), 345–358.
- [12] E. Dündar, U. Ulusu, F. Nuray, *On asymptotically ideal invariant equivalence of double sequences*, Thai J. Math. (in press).
- [13] A. Esi, and M. Açıkgöz, *On λ^2 -asymptotically double statistical equivalent sequences*, Int. J. Nonlinear Anal. Appl. **5** (2014), 16–21.
- [14] A. Esi, *On asymptotically double lacunary statistically equivalent sequences*, Appl. Math. Lett. **22** (2009), 1781–1785.
- [15] B. Hazarika, V. Kumar, *On Asymptotically double lacunary statistical equivalent sequences in ideal context*, J. Inequal. Appl. **2013**(543) (2013), DOI:10.1186/1029-242X-2013-543
- [16] B. Hazarika, *On asymptotically ideal equivalent sequences*, J. Egyptian Math. Soc. **23**(1) (2015), 67–72.
- [17] B. Hazarika, A. Alotaibi, S.A. Mohiuddine, *Statistical convergence inmeasure for double sequences of fuzzy-valued functions*, Soft Comput. **24** (2020), 6613–6622
- [18] U. Kadak, S.A. Mohiuddine, *Generalized statistically almost convergence based on the difference operator which includes the (p, q) -gamma function and related approximation theorems*, Results Math. (2018) 73:9.
- [19] E.E. Kara, M. Datan, M. İlkhani, *On lacunary ideal convergence of some sequences*, New Trends Math. Sci. **5**(1) (2017), 234–242.
- [20] V.A. Khan, N. Khan, *On some \mathcal{I} -convergent double sequence spaces defined by a modulus function*, Engineering, **5** (2013), 35–40.
- [21] G. Kılınç, İ. Solak, *Some double sequence spaces defined by a modulus function*, Gen. Math. Notes, **25**(2) (2014), 19–30.
- [22] P. Kostyrko, T. Šalát, W. Wilczyński, *\mathcal{I} -convergence*, Real Anal. Exchange, **26**(2) (2000), 669–686.
- [23] V. Kumar, A. Sharma, *Asymptotically lacunary equivalent sequences defined by ideals and modulus function*, Math. Sci. **6**(23) (2012), 5 pages.
- [24] V. Kumar, *On \mathcal{I} and \mathcal{I}^* -convergence of double sequences*, Math. Commun. **12** (2007), 171–181.
- [25] I.J. Maddox, *Sequence spaces defined by a modulus*, Math. Proc. Camb. Phil. Soc. **100** (1986), 161–166.
- [26] M. Marouf, *Asymptotic equivalence and summability*, Int. J. Math. Math. Sci. **16**(4) (1993), 755–762.

- [27] S.A. Mohiuddine, B.A.S. Alamri, *Generalization of equi-statistical convergence via weighted lacunary sequence with associated Korovkin and Voronovskaya type approximation theorems*, Rev. R. Acad. Cienc. Exactas Fis. Nat., Ser. A Mat. RACSAM, **113**(3) (2019), 1955–1973.
- [28] S.A. Mohiuddine, A. Asiri, B. Hazarika, *Weighted statistical convergence through difference operator of sequences of fuzzy numbers with application to fuzzy approximation theorems*, Int. J. Gen. Syst. **48**(5) (2019), 492–506.
- [29] S.A. Mohiuddine, B. Hazarika, M.A. Alghamdi, *Ideal relatively uniform convergence with Korovkin and Voronovskaya types approximation theorems*, Filomat, **33**(14) (2019), 4549–4560.
- [30] M. Mursaleen, *Matrix transformation between some new sequence spaces*, Houston J. Math. **9** (1983), 505–509.
- [31] M. Mursaleen, O. H. H. Edely, *Statistical convergence of double sequences*, J. Math. Anal. Appl. **288** (2003) 223–231.
- [32] M. Mursaleen, S.A. Mohiuddine, *On ideal convergence in probabilistic normed spaces*, Math. Slovaca, **62** (2012), 49–62.
- [33] H. Nakano, *Concave modulars*, J. Math. Soc. Japan, **5** (1953), 29–49.
- [34] N. Pancaroğlu Akin, *f -asymptotically lacunary ideal equivalence of double sequences*, J. Ineq. Appl. **2019** (2019), 1–11.
- [35] F. Nuray, H. Gök, U. Ulusu, \mathcal{I}_σ -convergence, Math. Commun. **16** (2011), 531–538.
- [36] F. Nuray, E. Dündar, U. Ulusu, *Wijsman \mathcal{I}_2 -convergence of double sequences of closed sets*, Pure Appl. Math. Lett. **2** (2014), 35–39.
- [37] N. Pancaroğlu, F. Nuray, *Statistical lacunary invariant summability*, Theoretical Math. Appl. **3**(2) (2013), 71–78.
- [38] N. Pancaroğlu, F. Nuray, *On invariant statistically convergence and lacunary invariant statistically convergence of sequences of sets*, Progress Appl. Math. **5**(2) (2013), 23–29.
- [39] N. Pancaroğlu, F. Nuray, *Invariant statistical convergence of sequences of sets with respect to a modulus function*, Abstr. Appl. Anal. **2014** (2014), 5 pages.
- [40] N. Pancaroğlu, F. Nuray, E. Savaş, *On asymptotically lacunary invariant statistical equivalent set sequences*, AIP Conf. Proc. **1558**(1) (2013), 780–781.
- [41] N. Pancaroğlu Akin, E. Dündar, U. Ulusu, *Wijsman lacunary \mathcal{I} -invariant convergence of sequences of sets*, Proc. Nat. Acad. Sci. India Sect. A. (2020), DOI:10.1007/s40010-020-00694-w
- [42] R.F. Patterson, *Rates of convergence for double sequences*, Southeast Asian Bull. Math. **26**(3) (2002), 469–478.
- [43] R.F. Patterson, E. Savaş, *Lacunary statistical convergence of double sequences*, Math. Commun. **10** (2005), 55–61.
- [44] R.F. Patterson, E. Savaş, *On asymptotically lacunary statistically equivalent sequences*, Thai J. Math. **4**(2) (2006), 267–272.
- [45] S. Pehlivan, B. Fisher, *Some sequences spaces defined by a modulus*, Math. Slovaca, **45** (1995), 275–280.
- [46] A. Pringsheim, *Zur theorie der zweifach unendlichen Zahlenfolgen*, Math. Ann. **53** (1900), 289–321.
- [47] R.A. Raimi, *Invariant means and invariant matrix methods of summability*, Duke Math. J. **30** (1963), 81–94.
- [48] E. Savaş, *Strong σ -convergent sequences*, Bull. Calcutta Math. **81** (1989), 295–300.
- [49] E. Savaş, F. Nuray, *On σ -statistically convergence and lacunary σ -statistically convergence*, Math. Slovaca, **43**(3) (1993), 309–315.
- [50] E. Savaş, *On \mathcal{I} -asymptotically lacunary statistical equivalent sequences*, Adv. Differ. Equ. **2013**(111) (2013), DOI:10.1186/1687-1847-2013-111
- [51] Y. Sever, U. Ulusu, E. Dündar, *On strongly \mathcal{I} and \mathcal{I}^* -lacunary convergence of sequences of sets*, AIP Conf. Proc. **1611**(1) (2014), 357–362.

- [52] Ş. Tortop, E. Dündar, *Wijsman \mathcal{I}_2 -invariant convergence of double sequences of sets*, J. Inequal. Spec. Funct. **9**(4) (2018), 90–100.
- [53] M.R. Türkmen, E. Dündar, *On lacunary statistical convergence of double sequences and some properties in fuzzy normed spaces*, J. Intell. Fuzzy Sys. **36**(2) (2019), 1683–1690, DOI: 10.3233/JIFS-18841
- [54] U. Ulusu, E. Dündar, *Asymptotically \mathcal{I}_2 -lacunary statistical equivalence of double sequences of sets*, J. Ineq. Special Func. **7**(2) (2016), 44–56.
- [55] U. Ulusu, E. Dündar, B. Aydın, *Asymptotically \mathcal{I}_2 -Cesaro equivalence of double sequences of sets*, J. Ineq. Special Func. **7**(4) (2016), 225–234.
- [56] U. Ulusu, E. Dündar, F. Nuray, *Lacunary \mathcal{I}_2 -invariant convergence and some properties*, Int. J. Anal. Appl. **16**(3) (2018), 317–327.
- [57] U. Ulusu, E. Dündar, *Asymptotically lacunary \mathcal{I}_2 -invariant equivalence*, J. Intell. Fuzzy Sys. **16**(1) (2019), 467–472.
- [58] S. Yegül, E. Dündar, *Statistical convergence of double sequences of functions and some properties in 2-normed spaces*, Facta Univ. Ser. Math. Inform. **33**(5) (2018), 705–719.

Erdoğan Dündar

Department of Mathematics,

Afyon Kocatepe University, 03200, Afyonkarahisar, Turkey

E-mail: edundar@aku.edu.tr

Nimet Pancaroğlu Akın

Department of Mathematics Education, Afyon Kocatepe University, 03200 Afyonkarahisar, Turkey

E-mail: npancaroglu@aku.edu.tr

Uğur Ulusu

Sivas Cumhuriyet University, 58140 Sivas, Turkey

E-mail: ugurulusu@cumhuriyet.edu.tr