

Kocatepe Veteriner Dergisi

ISSN: 1308-1594

e-ISSN: 2147-6853

Başlangıç: 2008

Yayımlanma Sıklığı: [Yılda 4 Sayı](#)

Yayıncı: Afyon Kocatepe Üniversitesi

Dergide makale ara



HAKKINDA

Alanında kaliteli makaleler yayımlamayı hedef edinmiş olan dergimizin veri tabanlarında taranmaya devam edebilmesi ve bir üst veri dizinlere katılabilmesi için, makale kabullerinde **Araştırma Makalesi, Olgu Sunumu ve Editöre Mektup** türünden yayınlarda **kaliteli ve İngilizce yazılmış yayınlara öncelik verilmesine; Derleme türünden yayınlarda ise davetli, spesifik alanlarda ve uluslararası okunabilirliği için sadece İngilizce olarak** kabul edilmesine karar verilmiştir.

"Mendeley" Kocatepe Veteriner Dergisi Kaynak Formatı için; View --> Citation Style --> More Styles --> Get More Styles --> Download Style

Enter URL kısmına link yapıştırın: "<http://csl.mendeley.com/styles/219358631/KVJ-en-son>"



182B

970B

20

223

+ Takip Et

21

ARŞİV

Son Sayılar

[2023 - Cilt: 16 Sayı: 1](#)
[2022 - Cilt: 15 Sayı: 4](#)
[2022 - Cilt: 15 Sayı: 3](#)
[2022 - Cilt: 15 Sayı: 2](#)

Erken Görünümdeki Sayılar

KEŞFET

[Makale Gönder](#)
[Hakemlik İsteği Gönder](#)
[Amaç ve Kapsam](#)
[Yazım Kuralları](#)
[Etik İlkeler ve Yayın Politikası](#)
[Ücret Politikası](#)
[Dergi Kurulları](#)
[İstatistikler](#)
[Dizinler](#)
[Tüm Sayılar](#)
[İletişim](#)

GÜNCEL SAYI

[2023 - Cilt: 16 Sayı: 1](#)
[Araştırma Makalesi](#)


[1. Emergent Contaminants in Freshwater Ecosystem: A case study from Turkey](#)

[Ayhan FİLAZİ](#) [Özgür KUZUKIRAN](#) [Gülçin AKCA](#) [Begüm YURDAKÖK DİKMEN](#) [Selin ÖZKAN KOTİLOĞLU*](#) [Mahmut SELVİ](#) [Figen ERKOÇ](#)

Sayfa: 1-15

Araştırma Makalesi

[2. Kronik enteropatili kedilerde bağırsak mikrobiyota analizleri ile kısa dönem rektal enema probiyotik sağaltımı](#)

[Kerem URAL](#) [Hasan ERDOĞAN](#) [Songül ERDOĞAN](#) [Serdar PAŞA](#) [Tuğba ASLAN](#) [Cansu BALIKÇI*](#) [Gamze GÖKÇAY](#)

Sayfa: 16-30

Araştırma Makalesi

[3. BUZAĞILARDA KLİNOPTİLOLİT \(ZEOLİT\) UYGULAMASININ ABOMAZAL BOŞALMA ORANINA ETKİSİ](#)

[Abdurrahman Kürşat ÖZSOY](#) [Tahir ÖZALP](#) [Hasan ERDOĞAN*](#)

Sayfa: 31-37

Araştırma Makalesi

[4. Gebeliğin Son Dönemindeki Holştayn Düvelerde Rasyona Rumen Korumalı Lizin Ve Metiyonin İlavesinin Bazı Metabolik Parametreler Üzerine Etkileri](#)

[Özge ÇAKMAK](#) [Hande Işıl AKBAĞ](#) [Canjir UYARLAR*](#)

Sayfa: 38-46

Araştırma Makalesi

[5. The Effects of In Ovo Administered Monosodium Glutamate on the Embryonic Development of Skeletal Muscle in Chickens](#)

[Ferhan BÖLÜKBAŞ*](#)

Sayfa: 47-56

Araştırma Makalesi

[6. Impact of Farm-Scale on Animal Management Practices in Pırlak Sheep Enterprises](#)

[Zehra BOZKURT*](#) [Özlem GÜCÜYENER HACAN](#) [Serdar KOÇAK](#) [Koray ÇELİKELİ](#) [Mustafa TEKERLİ](#) [Metin ERDOĞAN](#)

Sayfa: 57-69

Araştırma Makalesi

[7. Afyonkarahisar'da Tüketime Sunulan Ambalajlı ve Ambalajsız Dondurma Örneklerinin Mikrobiyolojik Kalite Parametrelerinin Araştırılması](#)

[Ulaş ACARÖZ*](#) [Recep KARA](#) [Zeki GÜRLER](#) [Ali SOYLU](#) [Nurtaç KÜÇÜKBÜĞRÜ](#)

Sayfa: 70-76

Araştırma Makalesi

[8. Differential Diagnostic Value of Serum Procalcitonin and Iron Levels in Diarrheic Neonatal Calves Caused by Escherichia coli and Rotavirus](#)

[Şükrü DEĞİRMENÇAY*](#) [Muhammed Sertaç EROĞLU](#) [Emre EREN](#)

Sayfa: 77-85

Araştırma Makalesi

[9. Geometric Morphometric Analysis of Skull in Honamlı, Hair, Kilis and Saanen Goat Using Dorsal Landmarks](#)

[Ahmet YAPRAK*](#) [Özcan ÖZGEL](#) [Yasin DEMİRASLAN](#)

Sayfa: 86-92

Araştırma Makalesi

[10. "Ruam Kâbil-i Tedavi midir? Ruama Karşı Aşı ve Serum İstihzar ve Tatbiki Mümkün müdür?" Başlıklı Ayrı Basımın Değerlendirilmesi Üzerine Bir Araştırma](#)[Abdullah KARA](#) [Erhan YÜKSEL](#) * [Özlem YÜKSEL](#) [Mustafa ÇAVDAROĞLU](#)

Sayfa: 93-108

Araştırma Makalesi

[11. Serum Levels of Mid-Regional Pro-adrenomedullin and Soluble Triggering Receptor Expressed on Myeloid Cells in Cattle with Pneumonia](#)[Mert SEZER](#) * [Enes AKYÜZ](#) [Oğuz MERHAN](#) [Uğur AYDIN](#) [Umut BATI](#) [Uğur YILDIZ](#) [Tahir GEZER](#) [Ekin Emre ERKILIÇ](#) [Ali Haydar KIRMIZIGÜL](#) [Gülbüz GÖKÇE](#)

Sayfa: 109-117

Olgu Sunumu

[1. Bir Boxer Irkı Köpekte Torakal Spondilozis Deformans Olgusu](#)[Zülfükar SARITAŞ](#) * [Zehra YAŞAR](#) [Fatma GÖRÜCÜ](#) [Yusuf KOÇ](#) [Musa KORKMAZ](#) [Musa YILMAZ](#)

Sayfa: 118-121

Olgu Sunumu


[2. A Case of Myiasis in a Biting Wound in a Red Fox \(*Vulpes vulpes*\)](#)[Zülfükar SARITAŞ](#) [Emine Hesna KANDIR](#) [Fatma GÖRÜCÜ](#) [Yusuf KOÇ](#) * [Beste SARAÇOĞLU](#)

Sayfa: 122-126

[13520](#) [13521](#) [13522](#) [13523](#) [14104](#)[14105](#) [14106](#) [14107](#) [14108](#)

Kocatepe Veteriner Dergisi

Dizinler

 Her veri tabanından dergilerin dizin güncelliği kontrol edilmelidir. DergiPark sorumluluk almamaktadır.

Atıf Dizinleri



TR Dizin

Diğer Dizinler



CAB Abstract



Türkiye Atıf Dizini



SOBIAD



Academindex

- Dergi Ana Sayfası
- Hakkında
- Amaç ve Kapsam
- Makale Gönder
- Dergi Kurulları
- İstatistikler
- Yazım Kuralları
- Etik İlkeler ve Yayın Politikası
- Ücret Politikası
- Dizinler
- Arşiv
- İletişim
- Mendeley

Kocatepe Veterinary Journal



Editör: Afyon Kocatepe Üniv. Veteriner Fak.

Yayın Formatı: Basılı ve Elektronik

Yayına Başladığı Yıl: 2008

Yıllık Yayın Sayısı: 4

Yayın Yapması Gereken Aylar: Mart, Haziran, Eylül, Aralık

Konu Alanları: Veterinerlik

Yayıncı: Afyon Kocatepe Üniv. Veteriner Fak.

Yayın Dili: Türkçe, İngilizce

Dizinlendiği Yıllar: 2015 - 2023 (Fen)

Konu Kategorisi: Fen > Veterinerlik

Makale Sayısı

448

Atıf Sayısı

223

Kendine Atıf Sayısı

41

Atıf Alan Makale Sayısı

126

Atıf Ortalaması

0,5

Kendine Atıf Oranı

%18,39

Makale & Atıf Sayısı



Makale Türleri



Kocatepe Veteriner Dergisi

Dergi Kurulları

Yayımcı/Sahibi

Prof. Dr. Turan CİVELEK

Dekan

Afyon Kocatepe Üniversitesi

Veteriner Fakültesi

Baş Editor

Prof. Dr. Musa KORKMAZ

Afyon Kocatepe Üniversitesi, Veteriner Fakültesi

Cerrahi Anabilim Dalı

musakorkmaz@aku.edu.tr

Editor

Doç. Dr. Recep KARA

Afyon Kocatepe Üniversitesi, Veteriner Fakültesi

Gıda Hijyeni ve Teknolojisi Anabilim Dalı

recepkala@aku.edu.tr

Doç. Dr. Deniz YENİ

Afyon Kocatepe Üniversitesi, Veteriner Fakültesi

Dölerme ve Suni Tohumlama Anabilim Dalı

dyeni@aku.edu.tr

Editör Yardımcıları

Doç. Dr. Muhammed Enes İNANÇ

Burdur Mehmet Akif Ersoy Üniversitesi, Veteriner Fakültesi

enesinanc@hotmail.com

Doç. Dr. Ruhi TÜRKMEN

Afyon Kocatepe Üniversitesi, Veteriner Fakültesi

ruhiturkmen@aku.edu.tr

[Dergi Ana Sayfası](#)

[Hakkında](#)

[Amaç ve Kapsam](#)

[Makale Gönder](#)

[Dergi Kurulları](#)

[İstatistikler](#)

[Yazım Kuralları](#)

[Etik İlkeler ve Yayın Politikası](#)

[Ücret Politikası](#)

[Dizinler](#)

[Arşiv](#)

[İletişim](#)

[Mendeley](#)

Dr. Öğr. Üyesi Özlem GÜCÜYENER HACAN

Afyon Kocatepe Üniversitesi, Veteriner Fakültesi

ogucuyener@gmail.com

Dr. Öğr. Üyesi Özlem ÖZDEN AKKAYA

Afyon Kocatepe Üniversitesi, Veteriner Fakültesi

ozlemozden@aku.edu.tr

Alan Editörleri**Prof. Dr. Alpaslan YILDIRIM**

Erciyes Üniversitesi, Veteriner Fakültesi

Parazitoloji Anabilim Dalı

yildirima@erciyes.edu.tr

Prof. Dr. Sadullah BAHAR

Selçuk Üniversitesi, Veteriner Fakültesi

Anatomi Anabilim Dalı

sbahar@selcuk.edu.tr

Prof. Dr. Akın YAKAN

Hatay Mustafa Kemal Üniversitesi, Veteriner Fakültesi

Genetik Anabilim Dalı

yakan@mku.edu.tr

Prof. Dr. Kemal Kaan TEKİNŞEN

Selçuk Üniversitesi, Veteriner Fakültesi

Gıda Hijyeni ve Teknolojisi Anabilim Dalı

kktekinsen@selcuk.edu.tr

Doç. Dr. Ali Evren HAYDARDEDEOĞLU

Aksaray Üniversitesi, Veteriner Fakültesi

İç Hastalıkları Anabilim Dalı

ahaydardedeoglu@aksaray.edu.tr

İstatistik Editörleri**Prof. Dr. İbrahim KILIÇ**

Afyon Kocatepe Üniversitesi, Veteriner Fakültesi

Biyoistatistik Anabilim Dalı

ibrahimkilig@aku.edu.tr

Doç. Dr. İlkay DOĞAN

Gaziantep Üniversitesi, Tıp Fakültesi

Biyoistatistik Anabilim Dalı

ilkaydogan@gantep.edu.tr

İngilizce Editörleri**Prof. Dr. İbrahim DEMİRKAN**

Afyon Kocatepe Üniversitesi, Veteriner Fakültesi

Cerrahi Anabilim Dalı

idemirkan@aku.edu.tr

Doç. Dr. Ulaş ACARÖZ

Afyon Kocatepe Üniversitesi, Veteriner Fakültesi

Gıda Hijyeni ve Teknolojisi Anabilim Dalı

ulasacaroz@aku.edu.tr

Organizasyon Komitesi**Prof. Dr. A. Fatih FİDAN**

Afyon Kocatepe Üniversitesi, Veteriner Fakültesi

Biyokimya Anabilim Dalı

ffidan@aku.edu.tr

Prof. Dr. Metin ERDOĞAN

Afyon Kocatepe Üniversitesi, Veteriner Fakültesi

Medikal Biyoloji ve Genetik Anabilim Dalı

erdogan@aku.edu.tr

Doç. Dr. Mustafa KABU

Afyon Kocatepe Üniversitesi, Veteriner Fakültesi

İç Hastalıkları Anabilim Dalı

mkabu@aku.edu.tr

Doç. Dr. Fatih AVDATEK

Afyon Kocatepe Üniversitesi, Veteriner Fakültesi

Dölerme ve Suni Tohumlama Anabilim Dalı

favdatek@aku.edu.tr

Dr. Ümit ÖZÇINAR

Afyon Kocatepe Üniversitesi, Veteriner Fakültesi

Hayvan Besleme ve Beslenme Hastalıkları Anabilim Dalı

umitozcinar@gmail.com

Dr. Barış DENK

Afyon Kocatepe Üniversitesi, Veteriner Fakültesi

Biyokimya Anabilim Dalı

bdenk@aku.edu.tr

Araş. Gör. Murat KIRIKKULAK

Afyon Kocatepe Üniversitesi, Veteriner Fakültesi

Dölerme ve Suni Tohumlama Anabilim Dalı

muratkirikkulak@gmail.com

Araş. Gör. Hülya ATİK

Afyon Kocatepe Üniversitesi, Veteriner Fakültesi

Fizyoloji Anabilim Dalı
hatik@aku.edu.tr

Araş. Gör. Beste SARAÇOĞLU

Afyon Kocatepe Üniversitesi, Veteriner Fakültesi
Yabani Hayvan Hast. ve Ekoloji Anabilim Dalı
bsaracoglu@aku.edu.tr

Araş. Gör. Eda DEMİRTAŞ

Afyon Kocatepe Üniversitesi, Veteriner Fakültesi
Medikal Biyoloji ve Genetik Anabilim Dalı
edemirtas@aku.edu.tr

13520 13521 13522 13523 14104

14105 14106 14107 14108

ERKEN GÖRÜNÜM



Cilt: 16 Sayı: 2

YIL: 2023

ARAŞTIRMA MAKALESİ

Araştırma Makalesi

1. Kütahya İlindeki Mezbahalarda Kesilen Sığırlarda Pasteurella Türlerinin İzolasyonu, İdentifikasyonu ve Antibiyotik Duyarlılıkları

Seydi Mehmet ARSLAN * Beytullah KENAR

Sayfa : 127-130

PDF

Araştırma Makalesi

2. Expression Patterns of Some Lipogenic Genes and Fatty Acid Profile of Liver in Food-Restricted Rats

Hüseyin ÖZKAN * İrem KARAASLAN Ufuk KAYA Sevdâ DALKIRAN İbrahim ALAKUŞ Baran ÇAMDEVİREN Hasan Hüseyin KEÇELİ Akın YAKAN

Sayfa : 131-142

PDF

Araştırma Makalesi

3. Effect of Anti-Mullerian Hormone, Metabolic Profile and Mineral Levels at Transition Period On The Calving – Conception Interval in Cows

Mustafa İLERİTÜRK * Özgür KAYNAR

Sayfa : 143-159

PDF

Araştırma Makalesi

4. The Effects of Restraint and Cold Restraint Stress on Coagulation Indicators in Wistar Albino Rats

Çağlasu KOÇ Mehmet EKİCİ *

Sayfa : 160-165

PDF

Araştırma Makalesi

5. Cryopreservation of Ram Semen Using Capsaicin Supplemented Tris Extender

Muhammed Enes İNANÇ Fatih AVDATEK Şükrü GÜNGÖR Mehmet Fuat GÜLHAN Kemal Tuna OLĞAÇ Barış DENK Deniz YENİ Umut TAŞDEMİR *

Sayfa : 166-173

PDF

Araştırma Makalesi

6. Serum C-Reactive Protein, Procalcitonin and Ceruloplasmin Concentrations in Dogs with Naturally Infected Ehrlichiosis

Gülten Emek TUNA * Gamze Sevrî EKREN AŞICI Pınar Alkim ULUTAŞ

Sayfa : 174-181

PDF

Araştırma Makalesi

7. Farklı Tırnak Lezyonu Bulunan Sığırlarda Serum ve Tırnak Dokusu Bazı İz Element Düzeylerinin Değerlendirilmesi

Burak KESGİN * Musa KORKMAZ

Sayfa : 182-194

PDF

OLGU SUNUMU

Olgu Sunumu

1. Deprem Sonrası Bir Safkan İngiliz Atta Gözlenen Stereotipik Davranış: Kendi Etrafında Dönme

Yavuzkan PAKSOY Özlem GÜCÜYENER HACAN *

PDF

Olgu Sunumu

2. The Efficiency of 7.2% Hypertonic Saline Solution on Echocardiographic Parameters in a Dog with Systemic Inflammatory Response Syndrome

Nilay ARSLAN * Cansu BALIKÇI Songül ERDOĞAN Hasan ERDOĞAN Kerem URAL

PDF

100

151

1

Arşiv

Cilt: 15 Sayı: 4, 31.12.2022

Cilt: 16 Sayı: 1, 31.03.2023

Cilt: 16 Sayı: 2



The Effects of Restraint and Cold Restraint Stress on Coagulation Indicators in Wistar Albino Rats

Çağlasu KOÇ¹, Mehmet EKİCİ^{2*}

¹3rd Year Student, Faculty of Veterinary Medicine, Sivas Cumhuriyet University, Sivas, Türkiye

²Department of Veterinary Physiology, Faculty of Veterinary Medicine, Sivas Cumhuriyet University, Sivas, Türkiye

ABSTRACT

The aim of this study was to investigate the effects of restraint and cold restraint stress from acute stress protocols on coagulation indicators in rats. The study was conducted in 18 male Wistar albino rats aged 8-10 weeks with a body weight of 180-220 g. After a one-week adaptation period, the rats were randomly divided into three groups (n=6/group). The animals in the control group were not exposed to any stress. Rats in the restraint group were housed in restrainers designed for rats, and their movement was restricted for 2 hours at room temperature. Rats in the cold restraint group were kept in the restrainer at +4°C for 2 hours. Blood samples were collected by cardiac puncture under ketamine and xylazine anesthesia in Vacutainer® tubes containing 3.2% sodium citrate (0.109 M trisodium citrate). Coagulation indicators (aPTT, PT, INR, fibrinogen, and D-dimer) were analyzed using an automated analyzer (Roche Cobas t511, Switzerland). Although acute stress (restraint and cold restraint stress) had no effect on aPTT and D-dimer levels (p>0.05), it increased PT and INR values (p<0.05) and decreased fibrinogen concentration (p<0.05). Consequently, acute stress may lead to platelet hypofunction in rats by prolonging PT, increasing INR, and decreasing fibrinogen concentration.

Keywords: Acute stress, aPTT, D-dimer, Fibrinogen, INR, PT

Kısıtlama ve Soğuk Kısıtlama Stresinin Wistar Albino Sıçanlarda Pıhtılaşma Göstergeleri Üzerindeki Etkileri

ÖZ

Bu çalışmanın amacı, kısıtlama ve soğuk kısıtlama stresi gibi iki farklı akut stres protokolünün sıçanlarda pıhtılaşma göstergeleri üzerindeki etkilerini araştırmaktır. Çalışma, vücut ağırlığı 180-220 g olan, 8-10 haftalık 18 erkek Wistar albino sıçan üzerinde gerçekleştirildi. Bir haftalık adaptasyon sürecinden sonra sıçanlar rastgele üç gruba (n=6/grup) ayrıldı. Kontrol grubundaki hayvanlar herhangi bir strese maruz bırakılmadı. Kısıtlama grubundaki ratlar, ratlar için tasarlanmış kısıtlayıcı içinde barındırıldı ve oda sıcaklığında 2 saat hareketleri kısıtlandı. Soğuk kısıtlama grubundaki ratlar +4°C'de 2 saat kısıtlayıcı içinde tutuldu. Kan numuneleri, %3,2 sodyum sitrat (0.109 M trisodyum sitrat) içeren Vacutainer® tüplerine ketamin ve ksilazin anestezisi altında kardiyak punksiyon yoluyla toplandı. Pıhtılaşma göstergeleri (aPTT, PT, INR, fibrinojen ve D-dimer) otomatik bir analiz cihazı (Roche Cobas t511, İsviçre) kullanılarak analiz edildi. Akut stres (kısıtlama ve soğuk kısıtlama stresi) aPTT ve D-dimer düzeylerine etki etmezken (p>0.05), PT ve INR değerlerini artırdı (p<0.05), fibrinojen konsantrasyonunu düşürdü (p<0.05). Sonuç olarak akut stres, PT'yi uzatarak, INR'yi artırarak ve fibrinojen konsantrasyonunu azaltarak sıçanlarda trombosit hipofonksiyonuna yol açabilir.

Anahtar kelimeler: Akut stres, aPTT, D-dimer, Fibrinojen, INR, PT

To cite this article: Koç Ç., Ekici M. The Effects of Restraint and Cold Restraint Stress on Coagulation Indicators in Wistar Albino Rats. Kocatepe Vet J. (2023) 16(2):160-165

Submission: 20.01.2023 Accepted: 16.05.2023 Published Online: 25.05.2023

ORCID ID; ÇK: 0000-0002-2925-6849, ME: 0000-0002-2163-6214

*Corresponding author e-mail: mehmetekici@cumhuriyet.edu.tr

INTRODUCTION

Stress-related factors such as increased platelet activation, endothelial dysfunction, coagulation, up-regulated inflammatory response, and altered fibrinolysis are crucial in the thrombotic processes linked to cardiovascular events (Sandrini et al. 2020). Numerous research on human hemostasis indicates that acute psychophysiological stress simultaneously stimulates coagulation and fibrinolysis, but also raises coagulation further, resulting in hypercoagulability (von Känel et al. 2001, Wirtz et al. 2008). In stressful conditions, the hypothalamic-pituitary-adrenal (HPA) axis and autonomic nerve system (ANS) are activated as a part of the organism's adaptive response, releasing glucocorticoids and catecholamines (norepinephrine and adrenaline), which either directly or indirectly affect hemostasis (Sandrini et al. 2020). Acute stress has different results on platelet reactivity in healthy subjects. Platelet surface glycoproteins, fibrinogen receptors, and P-selectin increased the expression of the organism to prevent excessive bleeding during acute stress, and the interaction with leukocytes and supports them come together (Strike et al. 2004). In addition to blood hyperviscosity, reduced plasma volume increased hematocrit, and activated coagulation factors, psychologically acute stress enhances plasma filtration in healthy individuals (Austin et al. 2012, Zraggen et al. 2005). On the other hand, a recent study found that the expression of P-selectin on the platelet surface was significantly increased in mice after 9 hours of stress (Pethaperumal et al. 2022). It has certainly been demonstrated in different animal models that sustained and intense stress increases platelet production and activation, enhancing the ability of thrombin and ADP to stimulate platelet aggregation and promote platelet-leukocyte interaction (Matsuhisa et al. 2014, Sandrini et al. 2017). It has been documented that individuals exposed to acute mental stress tend to increase the level of D-dimer with an increase in the level of some coagulation factors (FVII: C, FVIII: C, FXII: C, von Willebrand factor antigen (vWF)) and fibrinogen (Zraggen et al. 2005). In another study on humans, it was reported that aPTT decreased, and fibrinogen, FVIII: C, D-dimer, and PT values increased significantly in psychologically acute stress exposure (Austin et al. 2012). These inconsistencies may result from numerous study variations, including stress levels and platelet aggregation measuring techniques. In the literature, the effects of different stress protocols (repeated cold stress, acute vital stress, immobilization stress, water immersion stress, restraint stress, cold stress) on platelet functions have been investigated in previous rat and mouse stress studies (Bondarchuk et al. 2020, Hata et al. 1992, Hatu et al. 1991, Kawabata and Hata 1993, Loban'Chereda and Novosel'tseva 1990, Malyszko et al. 1994, Pethaperumal et al. 2022, Takeda et al. 1992).

In this study, we hypothesized that acute restraint and cold restraint stress protocols might have different effects on the coagulation profile and that mobility restriction with hypothermia might enhance the activation of the coagulation cascade. There is no experimental study on this subject in the literature. The aim of this study was to investigate the effects of restraint and cold restraint stress from acute stress protocols on coagulation indicators in rats.

MATERIAL AND METHODS

The study was carried out with approval from the Sivas Cumhuriyet University Animal Experiments Ethics Committee (Approval No: 65202830 - 050.04.04 - 640). The animals were provided by the Experimental Animals Application and Research Center of Sivas Cumhuriyet University. The animals were housed in standard care (appropriate ventilation, 12/12 h light/dark cycle, 21-23°C temperature, 35-60% humidity) and feeding (ad-libitum water and pelleted rat feed (DSA Agrifood Products Inc., Kırıkkale)) conditions. The content of pellet feed was as follows: Crude protein (24%), crude fiber (7.62%), crude oil (3%), crude ash (8.01%), phosphorus (0.75%), sodium (0.26%), and calcium (1.23%). The study was carried out on 18 male Wistar Albino rats, 8-10 weeks of age, with a body weight of 180-220 g. Rats were randomly divided into three groups (n=6/group) and underwent a one-week adaptation period to adapt to a new housing environment. Then, the rats were left quiescent for adaptation in the experimental room for 12 hours before applying the stress procedures.

Control group: Animals in this group did not be exposed to any stress. Blood samples were taken from animals simultaneously with acute stress group animals. Blood samples were taken from animals simultaneously with acute stress group animals. Blood samples were collected by cardiac puncture under ketamine (60 mg/kg, Keta-Control, Doğa İlaç, İstanbul) and xylazine (10 mg/kg, Ksilazol, Alivira, Ankara) anesthesia using 21-gauge needles, into 1.8 mL Vacutainer® tubes (Becton Dickinson, USA) with 3.2% sodium citrate (0.109 M trisodium citrate) and gently mixed six times. Blood was analyzed within 2 hours at the latest.

Restraint stress: Restraint group: In this stress protocol, rats were placed in restrictive cylinder-shaped plastics (Broome Restraint, 63.5 x 215.9mm (2.5" dia. x 8.5")) designed for the rat as previously described (Tu et al. 2019), and their movement was restricted for 2 hours at room temperature. Then, the procedure described in the control group was repeated.

Cold Restraint stress: Cold Restraint group: As previously described, rats were kept in the restrainer (Broome Restraint, 63.5 x 215.9mm (2.5" dia. x 8.5"))

at +4°C for 2 hours (Zhu et al. 2014). The procedure described in the control group was repeated. Coagulation indicators (aPTT, PT, and fibrinogen) and D-dimer were analyzed by optical coagulometric method and latex-based immunoturbidimetric method, respectively, using an analyzer (Roche Cobas t511, Switzerland) as previously described (Kitchen et al. 2018a, Kitchen et al. 2018b). Citrated blood tubes were centrifuged at 2500 g for 15 minutes. After centrifugation, samples were checked for preanalytical errors such as correct filling of the tube and the presence of hemolytic, lipemic, or icteric plasma that could affect test results. During the study, freshly collected plasma was stored in the sealed primary tube over the cell pellet. Roche aPTT Screen Reagent contains a mixture of purified phospholipids and silica particles as an activator that stimulates the formation of factor XIIIa. Calcium chloride is then added, which initiates the intrinsic coagulation cascade. Time from addition of calcium chloride to clot formation is measured. Roche PT Rec Reagent contains recombinant human thromboplastin with a heparin-neutralizing substance and calcium that, when added to citrated rat plasma, triggers activation of the extrinsic coagulation cascade. The time between addition of the reagent to plasma and formation of a fibrin clot is measured and expressed in seconds. The reagent lot-specific ISI is used to convert the patient's PT result in seconds to the international normalized ratio (INR) using the following formula $INR = (\text{patient PT} / \text{mean normal value PT})^{ISI}$. The ISI value for a given thromboplastin reagent is determined by a method comparison of the thromboplastin reagent to be standardized with an international reference thromboplastin. The fibrinogen test is a Clauss test using lyophilized bovine thrombin at a concentration

of 100 NIH units/ml with added stabilizers and buffers. The D-dimer test is a particle-enhanced immunoturbidimetric assay in which latex particles are coated with monoclonal antihuman D-dimer antibodies (mouse) at 0.12%. The start reagent is used together with a preservative/buffer solution at pH 8.2. Antigen/antibody complexes formed by the addition of samples containing D-dimer lead to an increase in the turbidity of the test reactants. The change in absorbance with time depends on the concentration of D-dimer epitopes in the sample. The aggregate is determined turbidimetrically.

Statistical Analysis

Data analysis was performed in GraphPad Prism 8.00 (GraphPad Software, San Diego, CA, United States). The normal distribution of the data was confirmed by the Shapiro-Wilk test. The one-way ANOVA test followed by Tukey's multiple comparisons test was used to evaluate differences in analytes. Results were expressed as mean \pm SD. Values of $p < 0.05$ were considered significant.

RESULTS

PT and INR values were statistically significant and higher in the restraint and cold restraint groups compared to the control group ($p < 0.05$). However, aPTT was not statistically significant between the groups ($p > 0.05$). Fibrinogen concentration was statistically significant and lower in the restraint group ($p < 0.001$) and cold restraint group compared to the control group ($p < 0.01$) (Figure 1). Interestingly, it yielded results < 0.2 in all animals in the D-dimer level groups. This value is within the reference range of 0-0.5 mg/L FEU (Table 1).

Table 1. D-Dimer concentration changes in groups.

Analyte	Groups			
	Control	Restraint	Cold Restraint	Reference range
D-Dimer (mg/L FEU)	<0.2	<0.2	<0.2	0-0.5




Figure 1: Coagulation parameters changes in rats exposed to restraint and cold restraint stress. * $p < 0.05$, ** $p < 0.01$, and *** $p < 0.001$ are statistically significant compared with control group according to one way ANOVA post hoc Tukey test. PT: Prothrombin time, INR: International normalized ratio, aPTT: Activated partial thromboplastin time.

DISCUSSION

In the present study, it was observed that two different acute stress protocols (restraint and cold restraint stress) had no clear effect on aPTT from coagulation indicators in rats, increasing PT and INR values, reducing fibrinogen concentration, had no effect on D-dimer concentration. To the best of the authors' knowledge, this study was the first to compare the effect of two acute stress protocols (restraint and cold restraint) on coagulation indicators in rats. According to the "glucocorticoid cascade hypothesis of aging" which explains the relationship between aging and stress response, immobilization stress increases corticosterone production in both aged and young animals, aged rats cannot reestablish their levels for at least four hours, indicating deficiencies in the negative feedback control of the HPA axis (Sapolsky et al. 1983, Sapolsky et al. 1986).

The use of young rats in our study is important in terms of the healthy response of the negative feedback mechanism to acute stress.

The temporal measurements of the extrinsic and intrinsic routes of coagulation are, respectively, PT and aPTT. Of the studies that examined at coagulation times in individuals, two of them found that there was no change in aPTT under acute psychological stress (de Boer et al. 2007, Zraggen et al. 2005), while the other reported a shortening of aPTT (von Känel et al. 2009). One research (von Känel et al. 2004) found no change in PT, another (de Boer et al. 2007) found a tendency toward shortening, and yet another (von Känel et al. 2009) found a considerable shortening. It has been documented that individuals exposed to acute mental stress tend to increase the level of D-dimer with an increase in the

level of some coagulation factors (FVII: C, FVIII: C, FXII: C, von Willebrand factor antigen (vWF)) and fibrinogen (Zraggen et al. 2005). In another study on humans, it was reported that aPTT decreased and fibrinogen, FVIII: C, D-dimer, and PT values increased significantly in psychologically acute stress exposure (Austin et al. 2012). The most recent study (von Känel et al. 2009) included prothrombotic alterations throughout two stress sessions, whereas earlier studies only looked at reactions at one stress session, which may help to explain why these contradictory results were obtained. More strong physiological effects are more likely to result from aggregating data from numerous assessments than from just one (Kamarck et al. 2000). It is unclear how acute psychological stress induces enhanced coagulation (von Känel et al. 2009, von Känel et al. 2001). Stress-induced catecholamine spillover and altered adrenergic receptor activity are two possible mechanisms.

Although there are considerable variations in blood coagulation when compared to the human standard, rat strains are commonly used in coagulation research (Lewis et al. 1985). The most notable distinction between human and rat blood coagulation is that rats have short clotting times that are pathological in human blood (García-Manzano et al. 2001). It has been reported that activated partial thromboplastin time (aPTT), thrombin time (TT) lengthened, prothrombin time (PT) did not change, and fibrinogen concentration decreased in rats exposed to repeated cold stress (Hatu et al. 1991). Another study reported inhibition of platelet aggregation, extrinsic coagulation, and a reduction in fibrinogen and antithrombin III levels in young rats subjected to experimental acute vital stress (prey animal; snake). In addition, increased platelet count with decreased aggregation capacity, hypocoagulation via the intrinsic pathway of plasma hemostasis activation, signs of thrombinemia, and increased antithrombin III have been reported in aged rats (Bondarchuk et al. 2020). While some studies have shown that platelet function increases response to acute stress (von Känel and Dimsdale 2000), it has been reported that it has no significant effect on platelet aggregation in mice (Matsuhisa et al. 2014). It has been shown that acute stress (water immersion restraint and cold restraint) in rats may cause a decrease in platelet aggregation (Malyszko et al. 1994, Takeda et al. 1992). In dogs under anesthesia, acute hypothermia led to transient platelet hypoaggregability (Yoshihara et al. 1985). In our study, although acute stress did not affect aPTT and D-dimer levels, it increased PT and INR values. In addition, fibrinogen concentration decreased under restraint and cold restraint stress. The possible reason for this is that platelet function may be impaired due to acute stress application, as previously stated in the rat study (Takeda et al. 1992). Platelet hypofunctions have been reported in the specific change in the rhythm of stressed (Hata et al. 1992, Kawabata and

Hata 1993) or immobilized (Loban²-Chereda and Novosel'tseva 1990) rats, and this is supported by our research. Another reason may be heparin which is secreted from the mast cells of rats under restraint stress (Umarova et al. 1997). Heparin increases blood anticoagulant potential in response to stress.

CONCLUSION

In conclusion, acute stress (restraint and cold restraint stress) in rats prolongs PT, increases INR, decreases fibrinogen concentration, and does not affect aPTT and D-dimer levels. Further studies on this topic are needed.

Financial Support: This study is the research result consisting of a part of the project accepted and supported within the scope of 2209-A University Students Research Projects Support Program.

Conflict of Interest: All authors have read and approved the study. There is no conflict of interest between the authors.

Authorship Contributions: The authors contributed equally to this study.

Ethics Committee Information: The study was carried out with approval from the Sivas Cumhuriyet University Animal Experiments Ethics Committee (Approval No: 65202830 - 050.04.04 - 640).

Acknowledgements: We appreciate Dr. Serkan Bolat's cooperation with the analysis of coagulation markers.

REFERENCES

- Austin AW, Wirtz PH, Patterson SM, Stutz M, von Känel R. Stress-induced alterations in coagulation: Assessment of a new hemoconcentration correction technique. *Psychosom Med.* 2012;74(3).
- de Boer D, Ring C, Wood M, Ford C, Jessney N, McIntyre D, Carroll D. Time course and mechanisms of mental stress-induced changes and their recovery: Hematocrit, colloid osmotic pressure, whole blood viscosity, coagulation times, and hemodynamic activity. *Psychophysiology.* 2007;44(4).
- Bondarchuk YA, Alekseeva O v., Shakhmatov II, Krinitsina IN. Response of the Hemostatic System to Acute Vital Stress in Rats. *Bull Exp Biol Med.* 2020;168(6).
- García-Manzano A, González-Llaven J, Lemini C, Rubio-Póo C. Standardization of rat blood clotting tests with reagents used for humans. *Proc West Pharmacol Soc.* 2001.
- Hata T, Kawabata A, Itoh E. Platelet hypoaggregability in rats exposed to SART stress (repeated cold stress). *Thromb Res.* 1992;65(4-5).

- Hatu T, Kawabata A, Itoh E.** Blood Coagulation and Fibrinolysis in SART-Stressed (Repeated Cold-Stressed) Rats and Drug Effects on the Altered Hemostatic Parameters. *The Japanese Journal of Pharmacology*. 1991;56(4).
- Kamarck TW, Debski TT, Manuck SB.** Enhancing the laboratory-to-life generalizability of cardiovascular reactivity using multiple occasions of measurement. *Psychophysiology*. 2000;37(4).
- von Känel R, Dimsdale JE.** Effects of sympathetic activation by adrenergic infusions on hemostasis in vivo. *Eur J Haematol*. 2000.
- von Känel R, Kudielka BM, Haeberli A, Stutz M, Fischer JE, Patterson SM.** Prothrombotic changes with acute psychological stress: Combined effect of hemoconcentration and genuine coagulation activation. *Thromb Res*. 2009;123(4).
- von Känel R, Mills PJ, Fainman C, Dimsdale JE.** Effects of psychological stress and psychiatric disorders on blood coagulation and fibrinolysis: A biobehavioral pathway to coronary artery disease? *Psychosom Med*. 2001;63(4).
- von Känel R, Preckel D, Zraggen L, Mischler K, Kudielka BM, Haeberli A, Fischer JE.** The effect of natural habituation on coagulation responses to acute mental stress and recovery in men. *Thromb Haemost*. 2004;92(6).
- Kawabata A, Hata T.** Characterization of platelet hypofunctions in rats under SART stress (repeated cold stress). *Thromb Res*. 1993;69(2).
- Kitchen S, Geisen U, Kappelmayer J, Quehenberger P, Lowe A, Jones R, Miles G, Boehm JG, Rozsnyai G.** Evaluating the analytical performance of four new coagulation assays for the measurement of fibrinogen, D-dimer and thrombin time. *Int J Lab Hematol*. 2018a;40(6).
- Kitchen S, Geisen U, Kappelmayer J, Quehenberger P, Drieß J, Lowe A, Jones R, Boehm JG, Miles G, Rozsnyai G.** Evaluating the analytical performance of five new coagulation assays for the measurement of prothrombin time and activated thromboplastin time. *Int J Lab Hematol*. 2018b Dec;40(6).
- Lewis JH, van Thiel DH, Hasiba U, Spero JA, Gavaler J.** Comparative hematology and coagulation: Studies on rodentia (Rats). *Comp Biochem Physiol A Physiol*. 1985;82(1).
- Loban'-Chereda GA, Novosel'tseva T v.** [Coagulative capacity of the blood and anti-aggregative activity of the vascular wall in rats subjected to immobilization stress]. *Fiziol Zh* [Internet]. 1990;36(2): 13—18. Available from: <http://europepmc.org/abstract/MED/2113873>
- Malyszko J, Urano T, Takada Y, Takada A.** Time-dependent changes in platelet aggregation, fibrinolytic activity, and peripheral serotonergic measures in rats subjected to water immersion restraint stress. *Haemostasis*. 1994;24(4).
- Matsuhisa F, Kitamura N, Satoh E.** Effects of acute and chronic psychological stress on platelet aggregation in mice. *Stress*. 2014;17(2).
- Pethaperumal S, Hung SC, Lien TS, Sun DS, Chang HH.** P-Selectin is a Critical Factor for Platelet-Mediated Protection on Restraint Stress-Induced Gastrointestinal Injury in Mice. *Int J Mol Sci*. MDPI; 2022;23(19).
- Sandrini L, Ieraci A, Amadio P, Popoli M, Tremoli E, Barbieri SS.** Apocynin Prevents Abnormal Megakaryopoiesis and Platelet Activation Induced by Chronic Stress. *Oxid Med Cell Longev*. 2017;2017.
- Sandrini L, Ieraci A, Amadio P, Zarà M, Barbieri SS.** Impact of acute and chronic stress on thrombosis in healthy individuals and cardiovascular disease patients. *Int J Mol Sci*. 2020.
- Sapolsky RM, Krey LC, McEwen BS.** The adrenocortical stress-response in the aged male rat: impairment of recovery from stress. *Exp Gerontol*. 1983;18(1).
- Sapolsky RM, Krey LC, McEwen BS.** The neuroendocrinology of stress and aging: the glucocorticoid cascade hypothesis. *Endocr Rev*. 1986; 7(3).
- Strike PC, Magid K, Brydon L, Edwards S, McEwan JR, Steptoe A.** Exaggerated platelet and hemodynamic reactivity to mental stress in men with coronary artery disease. *Psychosom Med*. 2004;66(4).
- Takeda H, Asaka M, Matsuno K, Ohtaki T, Miyazaki T.** Stress-induced gastric mucosal lesion and platelet aggregation in rats. *J Clin Gastroenterol*. 1992;14.
- Tu BX, Wang LF, Zhong XL, Hu ZL, Cao WY, Cui YH, Li SJ, Zou GJ, Liu Y, Zhou SF, Zhang WJ, Su JZ, Yan XX, Li F, Li CQ.** Acute restraint stress alters food-foraging behavior in rats: Taking the easier Way while suffered. *Brain Res Bull*. 2019;149.
- Umarova BA, Shapiro FB, Kogan AE, Strukova SM.** Mast cells secrete heparin during immobilization stress in rats. *Thromb Res*. 1997;85(3).
- Wirtz PH, Redwine LS, Baertschi C, Spillmann M, Ehlert U, von Känel AR.** Coagulation activity before and after acute psychosocial stress increases with age. *Psychosom Med*. 2008;70(4).
- Yoshihara H, Yamamoto T, Mihara H.** Changes in coagulation and fibrinolysis occurring in dogs during hypothermia. *Thromb Res*. 1985;37(4).
- Zraggen L, Fischer JE, Mischler K, Preckel D, Kudielka BM, von Känel R.** Relationship between hemoconcentration and blood coagulation responses to acute mental stress. *Thromb Res*. 2005;115(3).
- Zhu D, Tong Q, Liu W, Tian M, Xie W, Ji L, Shi J.** Angiotensin (1-7) protects against stress-induced gastric lesions in rats. *Biochem Pharmacol*. 2014;87(3).